



Rec'd PCT/PTO 22 DEC 2003  
PCT/GB 2003 / 002739



INVESTOR IN PEOPLE

## PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

The Patent Office  
Concept House  
Cardiff Road  
Newport  
South Wales  
NP10 8QQ

REC'D 22 JUL 2003  
WIPO PCT

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.


Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.

Signed

*Stephen Hordley*

Dated

11 July 2003



1/77

The Patent Office

Cardiff Road  
Newport  
South Wales  
NP10 8QQ

# Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

25JUN02 E728210-1 D00060  
P01/7700 0.00-0214528.2

1. Your reference

SMK/BP5956735

2. Patent application number

(The Patent Office will fill in this part)

0214528.2

24 JUN 2002

3. Full name, address and postcode of the or of each applicant (underline all surnames)

Patents ADP number (if you know it)

The University Court of The University of Aberdeen  
Regent Walk  
Aberdeen AB24 3FX  
UNITED KINGDOM

If the applicant is a corporate body, give the country/state of its incorporation

GB

4267126002

4. Title of the invention

Materials and Methods for Induction of Immune Tolerance

5. Name of your agent (if you have one)

MEWBURN ELLIS

Address for service in the United Kingdom to which all correspondence should be sent (including the postcode)

YORK HOUSE  
23 KINGSWAY  
LONDON  
WC2B 6HP

Patents ADP number (if you know it)

109006

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number

Country

Priority application number  
(if you know it)

Date of filing  
(day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing  
(day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request?

(Answer "Yes" if:

- a) any applicant named in part 3 is not an inventor, or
  - b) there is an inventor who is not named as an applicant, or
  - c) any named applicant is a corporate body.
- See note (d))

YES

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form 0

Description 48

Claim(s) 8

Abstract 0

Drawing(s) 20 + 20 En.

10. If you are also filing any of the following, state how many against each item

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for preliminary examination and search (Patents Form 9/77)

Request for substantive examination (Patents Form 10/77)

Any other documents (Please specify)

11.

I/We request the grant of a patent on the basis of this application.

Signature

*Simon M Kremer*

Date

24 June 2002

12. Name and daytime telephone number of person to contact in the United Kingdom

SIMON M KREMER

020 7240 4405

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

a) If you need help to fill in this form or you have any questions, please contact the Patent Office on 08459 500505.

b) Write your answers in capital letters using black ink or you may type them.

c) If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.

## Materials and methods for induction of immune tolerance

### Background to the invention

5 A wide variety of strategies are employed by viruses to evade  
immune mediated clearance, which can be considered to belong to one  
of three major mechanisms: escape, resistance and counterattack (Xu  
et al., 2001). Viruses escape immune recognition by disruption of  
antigen presentation pathways (Lorenzo et al., 2001) and epitope  
10 mutation (Erickson et al., 2001). Resistance is mediated by  
inhibiting apoptosis of virally infected cells. Counterattack  
comprises the killing of effector T cells (Mueller et al., 2001).  
Epstein-Barr virus (EBV) has been shown to avoid detection and  
clearance by such mechanisms. For example, escape of detection in  
Burkitt's lymphoma, where the cells express very low levels of MHC  
15 class I and adhesion molecules (Gregory et al., 1988), and Epstein-  
Barr Nuclear Antigen 1 escapes MHC class I presentation via its NH<sub>2</sub>  
terminal Gly/Ala repeat domain (Levitskaya et al., 1995, 1997).  
EBV also evades immune responses by resistance to apoptosis through  
its Bcl-2 homologue, BHRF-1 (Xu et al., 2001) and expression of the  
20 anti-apoptotic protein A20 induced by LMP1 (Fries et al., 1996).  
However, although EBV has been shown to avoid detection and  
clearance by these, and other mechanisms, it is still not clear how  
the virus maintains latent infection so successfully.

25 EBV is a human  $\gamma$ -herpes virus carried as a latent infection by more  
than 90% of adults, replicating in B-cells and nasopharyngeal  
epithelial cells (Kieff, 1996). The acute infection is controlled  
by a cytotoxic response predominantly against EBV Nuclear Antigens  
3A, 3B and 3C (Kieff, 1996), but, in all cases, the virus enters a  
30 latent state in B-cells (Kieff, 1996). LMP1 is part of a  
restricted panel of genes expressed during latency, and in several  
EBV-associated malignancies including Hodgkin's disease and  
nasopharyngeal carcinoma (Horikawa et al., 2000; Pallesen et al.,  
1991). The protein acts as a constitutively activated tumor  
35 necrosis factor receptor, transforming cells through activation of  
molecules including nuclear factor kappa B and the anti-apoptotic

protein A20 (Eliopoulos et al., 1996, 1997; Huen et al., 1995; Mosialos et al., 1995; Young et al., 1998).

The role of different CD4<sup>+</sup> T helper (Th) cell subsets in regulating the nature and efficacy of immune responses is increasingly recognized (Christensen et al., 2001; Groux et al., 1997; Levings and Roncarolo, 2000; Roncarolo et al., 2000, 2001; Shevach et al., 1998; Stephens and Mason, 2000; Thomsen et al., 2001). Initially, attention focused on mutual antagonism between Th1 and Th2 cells, which produce  $\gamma$ -interferon ( $\gamma$ -IFN) and IL-4 respectively (Mossman and Coffman, 1989), but further, T regulatory (Tr) cell subpopulations with important roles in immunoregulation and tolerance have now been defined (Groux et al., 1997; Levings and Roncarolo, 2000; Roncarolo et al., 2000, 2001; Shevach et al., 1998; Stephens and Mason, 2000). In particular, production of the Tr1 cytokine IL-10 can protect rodents against a number of immune-mediated diseases (Groux et al., 1997; Levings and Roncarolo, 2000), whilst Th3 cell secretion of transforming growth factor- $\beta$  prevents spontaneous autoimmunity (Gorelik and Flavell, 2000) and mediates some forms of oral tolerance (Weiner, 1997). Regulatory subpopulations characterized by CD25 expression have also been isolated from rodents (Seddon, and Mason, 2000; Shevach, 2000), and more recently from human peripheral blood (Jonuleit, et al., 2001; Levings, et al., 2001), but in most reports the suppressive effects of these cells are non-specific and not dependent on cytokine production. The importance of Tr cells in controlling immune-mediated disease raises the prospect that viruses may exploit such regulation as a fourth major mechanism to evade immune clearance.

Given that cells latently infected with EBV express LMP1, the question arises as to why this antigen fails to elicit protective cytotoxic immunity (Chapman et al., 2001; Khanna et al., 1998); cytotoxic T cells specific for LMP1 are notable for their absence from infected individuals (Chapman et al., 2001). Dukers et al. (2000) have recently suggested that LMP1 contains peptide motifs which can direct direct immunosuppressive effects on peripheral

Disclosure of the invention

5 The present inventors have found that certain infectious agents encode antigens comprising tolerogenic peptide sequences. By a "tolerogenic" peptide sequence is meant a sequence which, when administered to cells of the immune system, along with a target antigen, tolerises the cells to that target antigen.

10 Exposure to the tolerogenic sequence and the target antigen inhibits the capacity of the cells to mount an immune response to that target antigen on a subsequent challenge, regardless of whether or not the tolerogenic sequence is present for that subsequent challenge. However, although tolerised to the target  
15 antigen, populations of cells so treated retain their capacity to mount a response to other antigens in the absence of the tolerogenic sequence.

20 The types of immune response which can be inhibited in this way include "defensive" immune responses against foreign antigens, including those administered therapeutically, as well as "pathogenic" immune responses as seen in autoimmune and allergenic diseases. These responses are often characterised by lymphocyte proliferation, expression of cytokines such as IL-4 or gamma-IFN,  
25 and induction of antibody response.

The cells to be tolerised will be from an individual who has previously been infected with the infectious agent from which the tolerogenic peptide is derived.

30 Thus these tolerogenic sequences can induce antigen-specific tolerance of mononuclear leukocytes to target antigens. This activity therefore contrasts with the non-specific immunosuppressive effects attributed to some virus-derived  
35 peptides; e.g. from retroviral envelope proteins (Haraguchi et al., 1995) and EBV LMP1 protein (Dukers et al., 2000).

The present inventors have shown that it is possible to identify such sequences by testing their ability to induce expression of IL-10 in cells from a donor seropositive for the relevant infectious agent.

Accordingly, in a first aspect, the present invention provides a method for assessing the tolerogenicity of a test peptide sequence from an infectious agent, comprising the steps of:

- (i) contacting a cell population with said test peptide sequence,
- (ii) determining whether IL-10 expression in said cell population is increased, and optionally
- (iii) correlating the result of step (ii) with the tolerogenicity of the sequence,

wherein said cell population comprises mononuclear leukocytes from a donor previously infected by said infectious agent.

The term "mononuclear leukocytes" as used herein embraces T lymphocytes (including CD4<sup>+</sup> and CD8<sup>+</sup> T lymphocytes), B lymphocytes, natural killer (NK) cells, mononuclear phagocytes (monocytes and macrophages) and dendritic cells. Thus the cell population comprises one or more of these types of cells.

Preferably, the cell population comprises at least T lymphocytes, preferably CD4<sup>+</sup> lymphocytes, or at least one type of antigen presenting cell (APC). More preferably, the cell population comprises at least T lymphocytes, preferably CD4<sup>+</sup> lymphocytes, and at least one type of antigen presenting cell. An antigen presenting cell is any cell capable of presenting an antigen to a T lymphocyte in the context of an MHC class II molecule. Thus B lymphocytes, natural killer (NK) cells, mononuclear phagocytes

~~---(monocytes and macrophages) and dendritic cells are all considered~~  
to be APCs. However, the majority of nucleated cells are capable

exposed to pro-inflammatory cytokines, and so the cell population may further comprise APCs which would not normally be regarded as mononuclear leukocytes.

5 The cell population comprises mononuclear leukocytes derived from a donor previously infected by the relevant infectious agent. Preferably it can be demonstrated by an appropriate assay that the donor has previously raised an immune response against the  
10 infectious agent; for example, the donor may be seropositive for the infectious agent, i.e. have circulating antibodies specific for the infectious agent. Under some circumstances the donor may not have circulating antibodies specific for the infectious agent, for example where insufficient time has elapsed since infection for  
15 detectable levels of antibodies to be raised, or where a substantial time has elapsed since infection and antibody levels have fallen below the threshold of detectability. However, the term "seropositive" will be used throughout this specification to refer to any individual previously infected by the relevant  
20 infectious agent, regardless of actual serological status, and the term "seronegative" should be construed accordingly, i.e as referring to an individual not previously infected by the infectious agent.

25 The method may further comprise the steps of:

(i) (a) contacting a similar cell population from a donor not previously infected by said infectious agent with said test peptide sequence, and

30 (ii) (a) determining whether IL-10 expression in said cell population is increased,

and optionally

35 (ii) (b) comparing the results from step (ii) with the results from step (ii) (a).



In step (iii), the individual results, or any combination of the results, from any of steps (ii), (ii)(a) and (ii)(b) may be correlated with the tolerogenicity of the sequence. In general it is considered that the greater the level of IL-10 expression induced in the seropositive population by the test peptide, the more likely it is that the test peptide will be tolerogenic.

Whether or not IL-10 expression is increased may be determined by any appropriate method. Suitable methods include specific detection of IL-10 protein, e.g. by ELISA (Deveraux et al., 2000), flow cytometry (Kreft et al., 1992), non-competitive flow immunoassay (Kjellstrom et al., 2000), immunofluorescence (Scheffold et al., 2000) or immunoblot; by detection of IL-10 mRNA, e.g. by RT-PCR (Blaschke et al., 2000; Demay et al., 1996), or Northern blot; or by bioassay for IL-10 activity (Schlaak et al., 1994).

The present invention further provides a method for assessing the tolerogenicity of a test peptide sequence from an infectious agent towards a target antigen, comprising the steps of:

- (i) contacting a cell population with (a) said test peptide sequence and (b) a target antigen, to make a test composition, and
- (ii) re-contacting the cell population from said test composition with said target antigen.

wherein said cell population comprises mononuclear leukocytes from a donor previously infected by said infectious agent. Preferably, the cell population comprises at least one type of APC, which may or may not be a mononuclear leukocyte, as set out above.

In general the cell population will not be re-contacted with the test peptide in step (ii).

The method may further comprise the steps of:

(iii) assessing the response of said cell population to said target antigen, and optionally

5 (iv) correlating the result of step (iii) with the tolerogenicity of the test peptide sequence.

10 The response of the cell population to the second challenge with the target antigen may be assessed by any method that enables a tolerised population to be distinguished from a non-tolerised population. For example, a response of a non-tolerised population to a foreign antigen would be expected to include one or more of e.g. cell proliferation (typically lymphocyte proliferation), and expression of one or more cytokines (other than IL-10) such as IL-4, IL-2, IL-12 and gamma-IFN. Thus step (iii) may comprise the  
15 assessment of any one of these markers, or of any other suitable marker.

20 The method may be performed *in vivo* or *in vitro*. Preferably the method is performed *in vitro*, e.g. in culture. However the methods may be performed in any suitable model *in vivo*.

25 The purpose of re-contacting the cells with the target antigen in step (ii) is to confirm that the cells have been tolerised to the target antigen by the initial contact of step (i).

30 Therefore it is desirable that for step (ii), the test composition does not still contain appreciable amounts of the test peptide sequence, or of tolerogenic or immunosuppressive factors produced by the cells themselves, which might interfere with any reaction stimulated by the target antigen in step (ii). Therefore, the method may include the step of allowing the cells to rest between steps (i) and (ii), so that the activity of test peptide in the test composition is reduced, the cells are not still expressing tolerogenic factors which would interfere with any reaction in step  
35 (ii), and the activity of residual tolerogenic factors produced by the cells during or in response to the initial tolerogenic

challenge is reduced. IL-10 activity is used herein as a marker for tolerogenic factors generated by the PBMCs in step (i).

When performed *in vitro*, the method may additionally or alternatively comprise the step of washing the cells prior to step (ii). Washing may be performed in conventional fashion. Typically, the cells will be rested after washing. Fresh antigen presenting cells may be added before recontacting the cells with the target antigen in step (ii).

Without wishing to be bound by any particular theory, it is believed that IL-10 may play an effector role in inducing tolerance, so reduction of IL-10 activity may also be achieved by specific neutralisation, e.g. addition of a neutralising factor to the cells, such as a neutralising anti-IL-10 antibody.

The method may further comprise the step of contacting the cell population with a confirmatory antigen unrelated to the test sequence or the target antigen, to confirm that the cells retain their general reactive capability, even though their reactivity to the target antigen has been modified.

Any suitable antigen may be used as the target antigen or confirmatory antigen. These antigens may be primary antigens or recall antigens; that is to say, the cells in the assay may or may not have been exposed to them before. A typical primary antigen for assay use is KLH (keyhole limpet haemocyanin), while for donors previously immunised with Bacille Calmette-Guérin (BCG), purified protein derivative (PPD) from *Mycobacterium tuberculosis* is a suitable recall antigen. T cell mitogens such as Concanavalin A, which are generally regarded as relatively non-specific in their activation of T cells, can also be used as target or confirmatory antigens within the meaning of the present invention. It has been found that PBMCs can be rendered unresponsive to ConA by the techniques described herein, but still retain their ability to respond to other antigens.

A test peptide sequence which is capable of inducing IL-10 expression and/or antigen-specific tolerance in seropositive cells as described above may be regarded as a "tolerogenic peptide sequence".

5 A tolerogenic peptide sequence may therefore be used to modulate an immune response, either *in vivo* or *in vitro*, by administration to suitable seropositive mononuclear leukocytes along with a target antigen. This technique has a number of applications. For  
10 example, it may be used prophylactically, to prevent subsequent development of an inflammatory response to the target antigen, or to inhibit a pre-existing immune reaction to the target antigen.

15 Accordingly, in a further aspect, the present invention provides a method of tolerising a cell population to a target antigen, comprising contacting said cell population with

(a) a tolerogenic peptide sequence from an infectious agent,

20 and

(b) the target antigen,

25 wherein said cell population comprises mononuclear leukocytes from a donor seropositive for said infectious agent.

30 The cell population may be contacted with the tolerogenic peptide sequence and/or the target antigen directly. Alternatively, the cell population may be contacted with the tolerogenic peptide sequence and/or the target antigen indirectly, e.g. via APCs which would not normally be regarded as mononuclear leukocytes, as described above. Thus a population of APCs may be contacted with the tolerogenic peptide sequence and/or the target antigen, and the cell population subsequently contacted with the population of APCs.

35 The tolerogenic peptide and target antigen may be administered to the cell population, or to the population of APCs, either together

or separately, and in any order. Thus it is not intended that the tolerogenic peptide sequence and target antigen must necessarily be administered simultaneously.

5 Any or all of the steps described may be performed *in vitro*, *in vivo*, or *ex vivo*.

Thus all the steps described may be performed *in vitro*, e.g. in culture.

10

In some embodiments, a tolerogenic peptide sequence and a target antigen may be administered directly to a test subject or a subject to be treated, e.g. an individual who has previously been infected by the relevant infectious agent.

15

In alternative embodiments, a tolerogenic peptide sequence and a target antigen may be administered *in vitro* to a cell population comprising mononuclear leukocytes from such an individual. These cells may then be introduced into a test subject, or a subject to be treated, e.g. the subject from whom they were originally derived.

20

25

In alternative embodiments, a tolerogenic peptide sequence and a target antigen may be administered *in vitro* to a population of APCs. The population of APCs may then be contacted *in vitro* with a cell population comprising mononuclear leukocytes from an infected individual. That cell population, or a subset thereof e.g. some or all of the mononuclear leukocytes, may then be introduced into a test subject, or a subject to be treated, e.g. the subject from whom they were originally derived.

0

5

Alternatively, the population of APCs may be administered to a test subject, or a subject to be treated, e.g. the subject from whom they were originally derived. In this case contact between the cell population and the tolerogenic peptide sequence and target antigen takes place *in vivo* via the APCs.

Thus cells or tissues may be removed from a donor individual or individual to be treated, treated with the tolerogenic peptide sequence and a target antigen, and reintroduced to the donor. Suitable cells or tissues include particular type of antigen presenting cells, heterogeneous populations of cells, e.g. peripheral blood lymphocytes or subsets thereof, lymph nodes, etc.

Preferably, the cell population comprises at least T lymphocytes, preferably CD4<sup>+</sup> T lymphocytes. More preferably, the cell population comprises at least T lymphocytes, preferably CD4<sup>+</sup> T lymphocytes, and at least one type of APC. From the above description it can be seen that the cell population to be tolerised, may in some embodiments be considered to comprise cells *in situ* in a test subject or subject to be treated.

The test subject, or subject to be treated will typically be a mammal, and may be a human. In some embodiments, a test subject may be a non-human mammal e.g. a rodent, rabbit, etc. and will typically be seropositive for the infectious agent.

Certain infectious agents do not have animal models that are easy to manipulate. For example, the human pathogen EBV has no animal model. Therefore the test subject may be a non-human mammal with a severe combined immunodeficiency, comprising lymphocytes from a donor of the appropriate species seropositive for the infectious agent. By "severe combined immunodeficiency" is meant a defect in lymphocyte maturation, so that the affected animal has low or undetectable levels of mature T and/or B lymphocytes. The mammal may be a rodent, for example a mouse or rat, such as the SCID mouse. In preferred embodiments the non-human mammal with the severe combined immunodeficiency is reconstituted with human lymphocytes seropositive for EBV, e.g. from a seropositive donor. Suitable techniques are described in Mosier et al. (1988), McCune et al. (1988), Kamel-Reid et al. (1988), and Rowe et al. (1991). Similar techniques may be applied to create animal models of other conditions.

In any of the embodiments of the present invention, the target antigen may be a suitable test antigen as described above, or any antigen to which an inappropriate or undesirable immune response occurs or is likely to occur. Thus the target antigen may be one implicated in a disease state, e.g. a self antigen implicated in an autoimmune condition, such as rheumatoid arthritis, or an allergic state such as hayfever. The target antigen may be a protein, polypeptide or peptide, including an epitope of a protein, or any other suitable entity capable of provoking an immune reaction, such as polysaccharides, lipids, macromolecular complexes, cells, etc.

Examples of auto-immune diseases in which specific antigens have been identified as potentially pathogenically significant include multiple sclerosis (myelin basic protein), insulin-dependent diabetes mellitus (glutamic acid decarboxylase), insulin-resistant diabetes mellitus (insulin receptor), coeliac disease (gliadin), bullous pemphigoid (collagen type XVII), auto-immune haemolytic anaemia (Rh protein), auto-immune thrombocytopenia (GpIIb/IIIa), myasthenia gravis (acetylcholine receptor), Graves' disease (thyroid-stimulating hormone receptor), glomerulonephritis (collagen type IV), pernicious anaemia (intrinsic factor). Thus these antigens, or particular fragments or epitopes thereof may be suitable target antigens.

The target antigen may be an exogenous antigen which stimulates a response which also causes damage to host tissues. For example, acute rheumatic fever is caused by an antibody response to a Streptococcal antigen which cross-reacts with a cardiac muscle cell antigen. The target antigen may be one which provokes an atopic or allergic response, e.g. pollen (implicated in hayfever, e.g. Timothy Grass pollen), house dust mites (asthma), cosmetics, allergens administered via insect bites, nut allergens, or therapeutic products such as factor VIII, factor IX, blood group antigens, or monoclonal antibodies.

The methods of the present invention may be used to suppress

primary and secondary mixed lymphocyte reactions, graft rejection, and graft versus host disease. Thus a subject intended to receive a cellular transplant may be tolerised to antigens expressed by those cells. Alternatively, the transplant may be given in conjunction with tolerogenic peptide sequences as described herein, or nucleic acid encoding such peptide sequences, in order to tolerise the recipient to those cells. In preferred embodiments, some or all of the cells to be transplanted may be engineered to express tolerogenic peptides. Thus a cell to be transplanted may contain nucleic acid encoding a tolerogenic peptide sequence according to the present invention such that the cell is capable of expressing the tolerogenic peptide sequence. The optimum methodology will depend on the identity of the cells to be engineered. Antigen presenting cells, e.g. dendritic cells, etc., may be engineered to express the tolerogenic peptide sequence in such a manner that it is processed and presented in the context of the cells' own MHC II molecules. Other cell types may be engineered so that they secrete the expressed sequence, in order that it can be presented by neighbouring APCs.

In all of the aspects described herein, the infectious agent, from which the test or tolerogenic peptide sequence is derived, may be a virus. In preferred embodiments, the virus is a herpesvirus encoding a viral IL-10 homologue, preferably EBV.

The test or tolerogenic peptide sequence may be derived from an EBV protein, preferably EBV LMP1 protein or LMP2 protein. Thus the methods of the present invention extend to the use of LMP1 protein, LMP2 protein, or a portion or fragment thereof comprising a tolerogenic peptide sequence.

The test or tolerogenic peptide sequence may comprise one or more of the sequences p1 to p75, or p1' to p96'. If desired, more than one test or tolerogenic peptide sequence may be administered, either simultaneously or sequentially.



The present invention also provides a method of treating a disease mediated by an immune response against a target antigen, comprising the steps of administering (a) a tolerogenic peptide sequence from an infectious agent, and (b) the target antigen, to an individual seropositive for said infectious agent.

Nucleic acids encoding test or tolerogenic peptides, and/or target antigens, may be useful in all the methods of the present invention. As an alternative to administration of a peptide to cells, a nucleic acid encoding that peptide and capable of supporting its expression may be used instead. For example, DNA vaccination techniques are well known to the skilled person, as reviewed in Mor and Eliza (2001); Smith (2000); Schleef et al. (2000) and Apostolopoulos and Plebanski (2000). Thus where administration of a peptide sequence is referred to in any of the methods herein described, administration of a nucleic acid sequence encoding that peptide sequence is also envisaged. Thus contacting a cell population or population of antigen presenting cells with a peptide sequence is considered to encompass contacting the relevant cells with an appropriate nucleic acid.

Thus for example, the present invention further provides a method of tolerising a cell population to a target antigen, comprising contacting said cell population with

(a) a nucleic acid encoding said test peptide sequence, such that said test peptide sequence is expressed in said cell population, and

(b) the target antigen,

wherein said cell population comprises mononuclear leukocytes from a donor seropositive for said infectious agent.

Where the target antigen is a protein, polypeptide or peptide, a nucleic acid encoding the target antigen may be administered to the individual.

However this should not be taken to imply that the target antigen need necessarily be a protein, polypeptide or peptide.

5 Use of nucleic acids in this way is considered to be applicable, *mutatis mutandis*, to any corresponding embodiment of the present invention in which administration of a peptide sequence is referred to. When target antigens are protein or peptide, nucleic acids having appropriate coding sequences may likewise be administered instead. In related embodiments, cells may be contacted with  
10 peptides by contact with cells engineered to express the relevant peptides and either secrete them or present them in the context of MHC molecules.

15 The present invention further provides a pharmaceutical composition comprising a tolerogenic peptide sequence from an infectious agent and a target antigen, in admixture with a pharmaceutically acceptable carrier.

20 In preferred embodiments, the tolerogenic peptide sequence is derived from EBV, e.g. LMP1 or LMP2 as described above. Thus the composition may comprise EBV LMP1 protein, LMP2 protein, or a portion or fragment of either comprising a tolerogenic peptide sequence.

25 In preferred embodiments the tolerogenic peptide sequence may comprise one or more of the LMP1 peptide sequences P1 to P75, and/or one or more of the LMP2 peptide sequences P1' to P96' described herein.

30 The present invention further provides EBV LMP1 and LMP2 proteins, and portions or fragments of either, for example, the peptide sequences P1 to P75, or P1' to P96' comprising a tolerogenic peptide sequence, for use in a method of medical treatment.

35 The present invention further provides EBV LMP1 and LMP2 proteins, and portions or fragments thereof, for example, the peptide sequences P1 to P75, or P1' to P96' comprising a tolerogenic

peptide sequence, for use in the treatment of a condition mediated by an immune response directed against a target antigen.

The present invention further provides EBV LMP1 and LMP2 proteins, and portions or fragments thereof, for example, the LMP1 peptide sequences P1 to P75, and the LMP2 peptide sequences P1' to P96' comprising a tolerogenic peptide sequence, in the preparation of a medicament for the treatment of a condition mediated by an immune response directed against a target antigen. The medicament may further comprise the target antigen. The medicament will typically be formulated for administration to an individual previously infected by EBV.

In these and other aspects of the present invention, preferred peptides include P2, P4, P5, P6, P7, P8, P9, P10, P12, P13, P14, P15, P16, P17, P18, P20, P22, P23, P24, P25, P26, P27, P29, P30, P32, P34, P35, P39, P68, P71, P72. Particularly preferred peptides include P2, P4, P7, P14, P15, P18, P20, P22, P23, P24, and P32.

The condition may be, for example, type I diabetes mellitus, coeliac disease, multiple sclerosis, rheumatoid arthritis, systemic lupus erythematosus, myaesthesia gravis, autoimmune haemolytic anaemia and thrombocytopenia, an atopic response e.g. hay fever or asthma, or other allergy, e.g. to an allergen such as a pharmaceutical product or nut allergens, or an alloimmune response, e.g. graft rejection, graft versus host disease, or a response to therapeutic products such as factor VIII, or monoclonal antibody therapy. The target antigens described above may be useful for treatment of these conditions.

The compositions and medicaments described herein may comprise nucleic acids encoding tolerogenic peptides and/or target antigens, as appropriate.

Also provided is a pharmaceutical composition comprising a cell for treatment of a condition in a recipient in combination with a

acid encoding a tolerogenic peptide according to the present invention, such that said tolerogenic peptide sequence can be expressed by said cell.

- 5 The nucleic acid preferably encodes an EBV protein, e.g. LMP1 or LMP2, or a fragment thereof comprising a tolerogenic peptide sequence.

10 The tolerogenic peptide sequence and the target antigen may be administered together or separately. In preferred embodiments, they are administered together. They may be provided as an admixture of separate components, as a complex, or covalently associated. Where the target antigen is a protein, the tolerogenic peptide sequence and target antigen may be provided as a fusion  
15 protein. Use of fusion proteins in this manner is applicable to all aspects of the invention.

In any of the above-described aspects of the invention, the cell population to be tolerised may comprise mononuclear leukocytes from  
20 any suitable species. In preferred embodiments the mononuclear leukocytes are mammalian, e.g. from livestock animals such as horses, cattle, etc., from domestic animals, such as dogs, cats, etc., or from humans. Likewise, individuals to be treated by the methods of the present invention are preferably mammals, e.g.  
25 livestock animals such as horses, cattle, etc., domestic animals, such as dogs, cats, etc., and humans.

The term "peptide sequence" as used herein, whether a test or tolerogenic peptide sequence, should not be taken to refer solely  
30 to a free peptide consisting essentially or exclusively of that sequence, although this is encompassed by the present invention. Without wishing to be bound by any particular theory, it is believed that the methods of the present invention are effective as long as the relevant sequence can be presented to T cells by  
35 antigen presenting cells within the population. Thus it is believed that the test or tolerogenic peptide sequence may constitute a T cell epitope, in that it is capable of being

presented to T cells in the context of MHC molecules. Therefore the test or tolerogenic peptide sequence is preferably at least 6 amino acids in length, more preferably at least 8 amino acids in length.

Preferably the test or tolerogenic peptide sequence is capable of acting as an MHC class II-restricted T cell epitope. The chance that a peptide will be capable of acting as a T cell epitope can be determined by assessing its ability to bind to the antigen binding groove of MHC II molecules. Peptide motifs which bind particular MHC alleles are known, and computer programs are available which can identify such motifs within protein sequences (Sturniolo et al. (1999); Singh and Raghava (2001)).

The skilled person will be aware that any T cell that responds to a given peptide can also respond in a similar way to other peptides containing substitutions in residues that are not critical for MHC binding or T cell receptor recognition, and even to certain peptides that are substituted in critical residues. Such immunological cross reactivity of peptides can be demonstrated by showing that a particular T cell is capable of responding to more than one peptide. Such experiments may be performed using T cell clones. Techniques for cloning T cells are well known in the art. Without wishing to be bound by any particular theory, T cells of Tr1 phenotype may be implicated in the mechanism underlying the methods described herein. Such T cells do not proliferate significantly in response to stimulation, and suppress proliferation of other cells, and so can be difficult to clone. However, suitable techniques are known - see e.g. MacDonald et al. (2002).

Tolerogenic peptides derived from infectious agents described herein, or identified using the methods herein, may be used to screen for immunologically cross reactive peptides which exert similar tolerogenic effects by stimulating a similar or overlapping T cell population. Such cross reactive peptides may be considered

described herein. Thus the present invention provides a method for assessing the tolerogenicity of a test peptide sequence, comprising the steps of:

- 5 (i) contacting a first cell population with said test peptide sequence,
- (ii) contacting a second cell population with a control peptide sequence
- 10 (iii) determining whether IL-10 expression in each said cell population is increased, and optionally
- (iv) correlating the result of step (iii) with the tolerogenicity
- 15 of the test peptide sequence,

wherein each said cell population comprises mononuclear leukocytes from a donor previously infected by an infectious agent, and said control peptide sequence is derived from said infectious agent.

20 Thus typically, the control peptide sequence will have been previously shown to induce IL-10 expression in a cell population comprising mononuclear leukocytes from a donor previously infected by said infectious agent.

25 Preferably, the first and second cell populations are derived from the same donor individual. In preferred embodiments the first and second cell populations comprise T cell clones, preferably Tr1 T cell clones, shown to respond to the control peptide when appropriately presented by APCs.

30 The control peptide may comprise one or more of peptides P1 to P75 and/or P1' to P96' described herein.

35 The skilled person will also be aware that, because of the polymorphic nature of the MHC, most peptides will not be capable of binding to all MHC molecules. Thus compositions for use in the present invention may be tailored to a specific individual, by

selecting peptides likely to bind to their MHC. Alternatively, compositions may be designed to have a broader spectrum of activity, being applicable to a wider range of the population. This may be achieved by incorporating peptides capable of binding more than one MHC allele, and/or incorporating more than one test or tolerogenic peptide, each having different MHC specificity. These peptides may be provided in any appropriate form, e.g. as mixtures of separate peptides or as fusion proteins.

Therefore the test or tolerogenic peptide sequence may be administered as part of a longer peptide, polypeptide or protein. For example, the sequence may be used in the context of the whole or part of the full length native protein. The peptide, polypeptide or protein may be administered in any appropriate form, e.g. in native or denatured conformation.

It will be appreciated that any peptide, polypeptide or protein may comprise more than one tolerogenic peptide sequence within the meaning of the present invention. For example, the EBV LMP1 protein is believed to contain numerous individual peptide sequences capable of inducing tolerance to a target antigen in EBV-seropositive PBMCs, as described more fully in the Examples below.

Furthermore, a peptide, polypeptide or protein comprising one or more tolerogenic epitopes may be utilised in admixture with target antigen, or may, for example, be provided covalently coupled with a target antigen, either by chemical linkage, or, where the target antigen is a protein, as a fusion protein.

Peptides, polypeptides or proteins, including fusion proteins, for use in the methods or compositions of the present invention may be generated by any appropriate method, including chemical synthesis and recombinant expression.

The present invention further provides individual peptides having any one of the sequences of SEQ ID NO: 1 to 100, 101 to 200, 201 to 300, 301 to 400, 401 to 500, 501 to 600, 601 to 700, 701 to 800, 801 to 900, 901 to 1000, 1001 to 1100, 1101 to 1200, 1201 to 1300, 1301 to 1400, 1401 to 1500, 1501 to 1600, 1601 to 1700, 1701 to 1800, 1801 to 1900, 1901 to 2000, 2001 to 2100, 2101 to 2200, 2201 to 2300, 2301 to 2400, 2401 to 2500, 2501 to 2600, 2601 to 2700, 2701 to 2800, 2801 to 2900, 2901 to 3000, 3001 to 3100, 3101 to 3200, 3201 to 3300, 3301 to 3400, 3401 to 3500, 3501 to 3600, 3601 to 3700, 3701 to 3800, 3801 to 3900, 3901 to 4000, 4001 to 4100, 4101 to 4200, 4201 to 4300, 4301 to 4400, 4401 to 4500, 4501 to 4600, 4601 to 4700, 4701 to 4800, 4801 to 4900, 4901 to 5000, 5001 to 5100, 5101 to 5200, 5201 to 5300, 5301 to 5400, 5401 to 5500, 5501 to 5600, 5601 to 5700, 5701 to 5800, 5801 to 5900, 5901 to 6000, 6001 to 6100, 6101 to 6200, 6201 to 6300, 6301 to 6400, 6401 to 6500, 6501 to 6600, 6601 to 6700, 6701 to 6800, 6801 to 6900, 6901 to 7000, 7001 to 7100, 7101 to 7200, 7201 to 7300, 7301 to 7400, 7401 to 7500, 7501 to 7600, 7601 to 7700, 7701 to 7800, 7801 to 7900, 7901 to 8000, 8001 to 8100, 8101 to 8200, 8201 to 8300, 8301 to 8400, 8401 to 8500, 8501 to 8600, 8601 to 8700, 8701 to 8800, 8801 to 8900, 8901 to 9000, 9001 to 9100, 9101 to 9200, 9201 to 9300, 9301 to 9400, 9401 to 9500, 9501 to 9600, 9601 to 9700, 9701 to 9800, 9801 to 9900, 9901 to 10000, 10001 to 10100, 10101 to 10200, 10201 to 10300, 10301 to 10400, 10401 to 10500, 10501 to 10600, 10601 to 10700, 10701 to 10800, 10801 to 10900, 10901 to 11000, 11001 to 11100, 11101 to 11200, 11201 to 11300, 11301 to 11400, 11401 to 11500, 11501 to 11600, 11601 to 11700, 11701 to 11800, 11801 to 11900, 11901 to 12000, 12001 to 12100, 12101 to 12200, 12201 to 12300, 12301 to 12400, 12401 to 12500, 12501 to 12600, 12601 to 12700, 12701 to 12800, 12801 to 12900, 12901 to 13000, 13001 to 13100, 13101 to 13200, 13201 to 13300, 13301 to 13400, 13401 to 13500, 13501 to 13600, 13601 to 13700, 13701 to 13800, 13801 to 13900, 13901 to 14000, 14001 to 14100, 14101 to 14200, 14201 to 14300, 14301 to 14400, 14401 to 14500, 14501 to 14600, 14601 to 14700, 14701 to 14800, 14801 to 14900, 14901 to 15000, 15001 to 15100, 15101 to 15200, 15201 to 15300, 15301 to 15400, 15401 to 15500, 15501 to 15600, 15601 to 15700, 15701 to 15800, 15801 to 15900, 15901 to 16000, 16001 to 16100, 16101 to 16200, 16201 to 16300, 16301 to 16400, 16401 to 16500, 16501 to 16600, 16601 to 16700, 16701 to 16800, 16801 to 16900, 16901 to 17000, 17001 to 17100, 17101 to 17200, 17201 to 17300, 17301 to 17400, 17401 to 17500, 17501 to 17600, 17601 to 17700, 17701 to 17800, 17801 to 17900, 17901 to 18000, 18001 to 18100, 18101 to 18200, 18201 to 18300, 18301 to 18400, 18401 to 18500, 18501 to 18600, 18601 to 18700, 18701 to 18800, 18801 to 18900, 18901 to 19000, 19001 to 19100, 19101 to 19200, 19201 to 19300, 19301 to 19400, 19401 to 19500, 19501 to 19600, 19601 to 19700, 19701 to 19800, 19801 to 19900, 19901 to 20000, 20001 to 20100, 20101 to 20200, 20201 to 20300, 20301 to 20400, 20401 to 20500, 20501 to 20600, 20601 to 20700, 20701 to 20800, 20801 to 20900, 20901 to 21000, 21001 to 21100, 21101 to 21200, 21201 to 21300, 21301 to 21400, 21401 to 21500, 21501 to 21600, 21601 to 21700, 21701 to 21800, 21801 to 21900, 21901 to 22000, 22001 to 22100, 22101 to 22200, 22201 to 22300, 22301 to 22400, 22401 to 22500, 22501 to 22600, 22601 to 22700, 22701 to 22800, 22801 to 22900, 22901 to 23000, 23001 to 23100, 23101 to 23200, 23201 to 23300, 23301 to 23400, 23401 to 23500, 23501 to 23600, 23601 to 23700, 23701 to 23800, 23801 to 23900, 23901 to 24000, 24001 to 24100, 24101 to 24200, 24201 to 24300, 24301 to 24400, 24401 to 24500, 24501 to 24600, 24601 to 24700, 24701 to 24800, 24801 to 24900, 24901 to 25000, 25001 to 25100, 25101 to 25200, 25201 to 25300, 25301 to 25400, 25401 to 25500, 25501 to 25600, 25601 to 25700, 25701 to 25800, 25801 to 25900, 25901 to 26000, 26001 to 26100, 26101 to 26200, 26201 to 26300, 26301 to 26400, 26401 to 26500, 26501 to 26600, 26601 to 26700, 26701 to 26800, 26801 to 26900, 26901 to 27000, 27001 to 27100, 27101 to 27200, 27201 to 27300, 27301 to 27400, 27401 to 27500, 27501 to 27600, 27601 to 27700, 27701 to 27800, 27801 to 27900, 27901 to 28000, 28001 to 28100, 28101 to 28200, 28201 to 28300, 28301 to 28400, 28401 to 28500, 28501 to 28600, 28601 to 28700, 28701 to 28800, 28801 to 28900, 28901 to 29000, 29001 to 29100, 29101 to 29200, 29201 to 29300, 29301 to 29400, 29401 to 29500, 29501 to 29600, 29601 to 29700, 29701 to 29800, 29801 to 29900, 29901 to 30000, 30001 to 30100, 30101 to 30200, 30201 to 30300, 30301 to 30400, 30401 to 30500, 30501 to 30600, 30601 to 30700, 30701 to 30800, 30801 to 30900, 30901 to 31000, 31001 to 31100, 31101 to 31200, 31201 to 31300, 31301 to 31400, 31401 to 31500, 31501 to 31600, 31601 to 31700, 31701 to 31800, 31801 to 31900, 31901 to 32000, 32001 to 32100, 32101 to 32200, 32201 to 32300, 32301 to 32400, 32401 to 32500, 32501 to 32600, 32601 to 32700, 32701 to 32800, 32801 to 32900, 32901 to 33000, 33001 to 33100, 33101 to 33200, 33201 to 33300, 33301 to 33400, 33401 to 33500, 33501 to 33600, 33601 to 33700, 33701 to 33800, 33801 to 33900, 33901 to 34000, 34001 to 34100, 34101 to 34200, 34201 to 34300, 34301 to 34400, 34401 to 34500, 34501 to 34600, 34601 to 34700, 34701 to 34800, 34801 to 34900, 34901 to 35000, 35001 to 35100, 35101 to 35200, 35201 to 35300, 35301 to 35400, 35401 to 35500, 35501 to 35600, 35601 to 35700, 35701 to 35800, 35801 to 35900, 35901 to 36000, 36001 to 36100, 36101 to 36200, 36201 to 36300, 36301 to 36400, 36401 to 36500, 36501 to 36600, 36601 to 36700, 36701 to 36800, 36801 to 36900, 36901 to 37000, 37001 to 37100, 37101 to 37200, 37201 to 37300, 37301 to 37400, 37401 to 37500, 37501 to 37600, 37601 to 37700, 37701 to 37800, 37801 to 37900, 37901 to 38000, 38001 to 38100, 38101 to 38200, 38201 to 38300, 38301 to 38400, 38401 to 38500, 38501 to 38600, 38601 to 38700, 38701 to 38800, 38801 to 38900, 38901 to 39000, 39001 to 39100, 39101 to 39200, 39201 to 39300, 39301 to 39400, 39401 to 39500, 39501 to 39600, 39601 to 39700, 39701 to 39800, 39801 to 39900, 39901 to 40000, 40001 to 40100, 40101 to 40200, 40201 to 40300, 40301 to 40400, 40401 to 40500, 40501 to 40600, 40601 to 40700, 40701 to 40800, 40801 to 40900, 40901 to 41000, 41001 to 41100, 41101 to 41200, 41201 to 41300, 41301 to 41400, 41401 to 41500, 41501 to 41600, 41601 to 41700, 41701 to 41800, 41801 to 41900, 41901 to 42000, 42001 to 42100, 42101 to 42200, 42201 to 42300, 42301 to 42400, 42401 to 42500, 42501 to 42600, 42601 to 42700, 42701 to 42800, 42801 to 42900, 42901 to 43000, 43001 to 43100, 43101 to 43200, 43201 to 43300, 43301 to 43400, 43401 to 43500, 43501 to 43600, 43601 to 43700, 43701 to 43800, 43801 to 43900, 43901 to 44000, 44001 to 44100, 44101 to 44200, 44201 to 44300, 44301 to 44400, 44401 to 44500, 44501 to 44600, 44601 to 44700, 44701 to 44800, 44801 to 44900, 44901 to 45000, 45001 to 45100, 45101 to 45200, 45201 to 45300, 45301 to 45400, 45401 to 45500, 45501 to 45600, 45601 to 45700, 45701 to 45800, 45801 to 45900, 45901 to 46000, 46001 to 46100, 46101 to 46200, 46201 to 46300, 46301 to 46400, 46401 to 46500, 46501 to 46600, 46601 to 46700, 46701 to 46800, 46801 to 46900, 46901 to 47000, 47001 to 47100, 47101 to 47200, 47201 to 47300, 47301 to 47400, 47401 to 47500, 47501 to 47600, 47601 to 47700, 47701 to 47800, 47801 to 47900, 47901 to 48000, 48001 to 48100, 48101 to 48200, 48201 to 48300, 48301 to 48400, 48401 to 48500, 48501 to 48600, 48601 to 48700, 48701 to 48800, 48801 to 48900, 48901 to 49000, 49001 to 49100, 49101 to 49200, 49201 to 49300, 49301 to 49400, 49401 to 49500, 49501 to 49600, 49601 to 49700, 49701 to 49800, 49801 to 49900, 49901 to 50000, 50001 to 50100, 50101 to 50200, 50201 to 50300, 50301 to 50400, 50401 to 50500, 50501 to 50600, 50601 to 50700, 50701 to 50800, 50801 to 50900, 50901 to 51000, 51001 to 51100, 51101 to 51200, 51201 to 51300, 51301 to 51400, 51401 to 51500, 51501 to 51600, 51601 to 51700, 51701 to 51800, 51801 to 51900, 51901 to 52000, 52001 to 52100, 52101 to 52200, 52201 to 52300, 52301 to 52400, 52401 to 52500, 52501 to 52600, 52601 to 52700, 52701 to 52800, 52801 to 52900, 52901 to 53000, 53001 to 53100, 53101 to 53200, 53201 to 53300, 53301 to 53400, 53401 to 53500, 53501 to 53600, 53601 to 53700, 53701 to 53800, 53801 to 53900, 53901 to 54000, 54001 to 54100, 54101 to 54200, 54201 to 54300, 54301 to 54400, 54401 to 54500, 54501 to 54600, 54601 to 54700, 54701 to 54800, 54801 to 54900, 54901 to 55000, 55001 to 55100, 55101 to 55200, 55201 to 55300, 55301 to 55400, 55401 to 55500, 55501 to 55600, 55601 to 55700, 55701 to 55800, 55801 to 55900, 55901 to 56000, 56001 to 56100, 56101 to 56200, 56201 to 56300, 56301 to 56400, 56401 to 56500, 56501 to 56600, 56601 to 56700, 56701 to 56800, 56801 to 56900, 56901 to 57000, 57001 to 57100, 57101 to 57200, 57201 to 57300, 57301 to 57400, 57401 to 57500, 57501 to 57600, 57601 to 57700, 57701 to 57800, 57801 to 57900, 57901 to 58000, 58001 to 58100, 58101 to 58200, 58201 to 58300, 58301 to 58400, 58401 to 58500, 58501 to 58600, 58601 to 58700, 58701 to 58800, 58801 to 58900, 58901 to 59000, 59001 to 59100, 59101 to 59200, 59201 to 59300, 59301 to 59400, 59401 to 59500, 59501 to 59600, 59601 to 59700, 59701 to 59800, 59801 to 59900, 59901 to 60000, 60001 to 60100, 60101 to 60200, 60201 to 60300, 60301 to 60400, 60401 to 60500, 60501 to 60600, 60601 to 60700, 60701 to 60800, 60801 to 60900, 60901 to 61000, 61001 to 61100, 61101 to 61200, 61201 to 61300, 61301 to 61400, 61401 to 61500, 61501 to 61600, 61601 to 61700, 61701 to 61800, 61801 to 61900, 61901 to 62000, 62001 to 62100, 62101 to 62200, 62201 to 62300, 62301 to 62400, 62401 to 62500, 62501 to 62600, 62601 to 62700, 62701 to 62800, 62801 to 62900, 62901 to 63000, 63001 to 63100, 63101 to 63200, 63201 to 63300, 63301 to 63400, 63401 to 63500, 63501 to 63600, 63601 to 63700, 63701 to 63800, 63801 to 63900, 63901 to 64000, 64001 to 64100, 64101 to 64200, 64201 to 64300, 64301 to 64400, 64401 to 64500, 64501 to 64600, 64601 to 64700, 64701 to 64800, 64801 to 64900, 64901 to 65000, 65001 to 65100, 65101 to 65200, 65201 to 65300, 65301 to 65400, 65401 to 65500, 65501 to 65600, 65601 to 65700, 65701 to 65800, 65801 to 65900, 65901 to 66000, 66001 to 66100, 66101 to 66200, 66201 to 66300, 66301 to 66400, 66401 to 66500, 66501 to 66600, 66601 to 66700, 66701 to 66800, 66801 to 66900, 66901 to 67000, 67001 to 67100, 67101 to 67200, 67201 to 67300, 67301 to 67400, 67401 to 67500, 67501 to 67600, 67601 to 67700, 67701 to 67800, 67801 to 67900, 67901 to 68000, 68001 to 68100, 68101 to 68200, 68201 to 68300, 68301 to 68400, 68401 to 68500, 68501 to 68600, 68601 to 68700, 68701 to 68800, 68801 to 68900, 68901 to 69000, 69001 to 69100, 69101 to 69200, 69201 to 69300, 69301 to 69400, 69401 to 69500, 69501 to 69600, 69601 to 69700, 69701 to 69800, 69801 to 69900, 69901 to 70000, 70001 to 70100, 70101 to 70200, 70201 to 70300, 70301 to 70400, 70401 to 70500, 70501 to 70600, 70601 to 70700, 70701 to 70800, 70801 to 70900, 70901 to 71000, 71001 to 71100, 71101 to 71200, 71201 to 71300, 71301 to 71400, 71401 to 71500, 71501 to 71600, 71601 to 71700, 71701 to 71800, 71801 to 71900, 71901 to 72000, 72001 to 72100, 72101 to 72200, 72201 to 72300, 72301 to 72400, 72401 to 72500, 72501 to 72600, 72601 to 72700, 72701 to 72800, 72801 to 72900, 72901 to 73000, 73001 to 73100, 73101 to 73200, 73201 to 73300, 73301 to 73400, 73401 to 73500, 73501 to 73600, 73601 to 73700, 73701 to 73800, 73801 to 73900, 73901 to 74000, 74001 to 74100, 74101 to 74200, 74201 to 74300, 74301 to 74400, 74401 to 74500, 74501 to 74600, 74601 to 74700, 74701 to 74800, 74801 to 74900, 74901 to 75000, 75001 to 75100, 75101 to 75200, 75201 to 75300, 75301 to 75400, 75401 to 75500, 75501 to 75600, 75601 to 75700, 75701 to 75800, 75801 to 75900, 75901 to 76000, 76001 to 76100, 76101 to 76200, 76201 to 76300, 76301 to 76400, 76401 to 76500, 76501 to 76600, 76601 to 76700, 76701 to 76800, 76801 to 76900, 76901 to 77000, 77001 to 77100, 77101 to 77200, 77201 to 77300, 77301 to 77400, 77401 to 77500, 77501 to 77600, 77601 to 77700, 77701 to 77800, 77801 to 77900, 77901 to 78000, 78001 to 78100, 78101 to 78200, 78201 to 78300, 78301 to 78400, 78401 to 78500, 78501 to 78600, 78601 to 78700, 78701 to 78800, 78801 to 78900, 78901 to 79000, 79001 to 79100, 79101 to 79200, 79201 to 79300, 79301 to 79400, 79401 to 79500, 79501 to 79600, 79601 to 79700, 79701 to 79800, 79801 to 79900, 79901 to 80000, 80001 to 80100, 80101 to 80200, 80201 to 80300, 80301 to 80400, 80401 to 80500, 80501 to 80600, 80601 to 80700, 80701 to 80800, 80801 to 80900, 80901 to 81000, 81001 to 81100, 81101 to 81200, 81201 to 81300, 81301 to 81400, 81401 to 81500, 81501 to 81600, 81601 to 81700, 81701 to 81800, 81801 to 81900, 81901 to 82000, 82001 to 82100, 82101 to 82200, 82201 to 82300, 82301 to 82400, 82401 to 82500, 82501 to 82600, 82601 to 82700, 82701 to 82800, 82801 to 82900, 82901 to 83000, 83001 to 83100, 83101 to 83200, 83201 to 83300, 83301 to 83400, 83401 to 83500, 83501 to 83600, 83601 to 83700, 83701 to 83800, 83801 to 83900, 83901 to 84000, 84001 to 84100, 84101 to 84200, 84201 to 84300, 84301 to 84400, 84401 to 84500, 84501 to 84600, 84601 to 84700, 84701 to 84800, 84801 to 84900, 84901 to 85000, 85001 to 85100, 85101 to 85200, 85201 to 85300, 85301 to 85400, 85401 to 85500, 85501 to 85600, 85601 to 85700,

P7, P8, P9, P10, P12, P13, P14, P15, P16, P17, P18, P20, P22, P23, P24, P25, P26, P27, P29, P30, P32, P34, P35, P39, P68, P71, P72. Particularly preferred peptides have sequences of P2, P4, P7, P14, P15, P18, P20, P22, P23, P24, and P32.

Thus in a further aspect, the present invention provides isolated nucleic acid molecules encoding the test and tolerogenic sequences of the present invention. The open reading frame may be contiguous with an open reading frame encoding a desired target antigen, in order to encode a fusion protein as described above.

In further aspects, the present invention provides an expression vector comprising the above tolerogenic sequence-encoding nucleic acid, operably linked to control sequences to direct its expression, as well as host cells transformed with the vectors. The present invention also includes a method of producing peptides of the preceding aspect, comprising culturing the host cells and isolating the tolerogenic peptides thus produced.

In order to obtain expression of nucleic acids encoding test, tolerogenic or target antigen sequences, the sequences can be incorporated into a vector having control sequences operably linked to the encoding nucleic acid to control its expression. The vectors may include other sequences such as promoters or enhancers to drive the expression of the inserted nucleic acid, nucleic acid sequences so that the tolerogenic sequence peptide is produced as a fusion, e.g. with one or more other such tolerogenic sequences, or with one or more target antigens, and/or nucleic acid encoding secretion signals so that the peptide produced in the host cell is secreted from the cell. Peptides/polypeptides/proteins can then be obtained by transforming the vectors into host cells in which the vector is functional, culturing the host cells so that the peptide is produced and recovering the peptide from the host cells or the surrounding medium. Prokaryotic and eukaryotic cells are used for this purpose in the art, including strains of *E. coli*, yeast, and eukaryotic cells such as COS or CHO cells.



Suitable vectors can be chosen or constructed, containing appropriate regulatory sequences, including promoter sequences, terminator fragments, polyadenylation sequences, enhancer sequences, marker genes and other sequences as appropriate.

Vectors may be plasmids, viral e.g. 'phage, or phagemid, as appropriate. For further details see, for example, "Molecular Cloning: a Laboratory Manual": 2nd edition, Sambrook et al., 1989, Cold Spring Harbor Laboratory Press.

Cells and techniques may be selected such as to permit or enhance the folding and/or formation of disulphide bridges (see e.g. "Protein Folding" by R. Hermann, Pub. 1993, European Patent Office, The Hague, Netherlands, ISBN 90-9006173-8).

Peptides may be synthesized by any suitable method, such as by exclusively solid-phase techniques, by partial solid-phase techniques, by fragment condensation or by classical solution couplings. In conventional solution phase peptide synthesis, the peptide chain can be prepared by a series of coupling reactions in which the constituent amino acids are added to the growing peptide chain in the desired sequence.

Briefly, N-alpha-protected amino acid anhydrides are prepared in crystallized form or prepared freshly in solution and used for successive amino acid addition at the N-terminus. At each residue addition, the growing peptide (on a solid support) is acid treated to remove the N-alpha-protective group, washed several times to remove residual acid and to promote accessibility of the peptide terminus to the reaction medium. The peptide is then reacted with an activated N-protected amino acid symmetrical anhydride, and the solid support is washed. At each residue-addition step, the amino acid addition reaction may be repeated for a total of two or three separate addition reactions, to increase the percent of growing peptide molecules which are reacted. Typically, 1-2 reaction cycles are used for the first twelve residue additions, and 2-3 reaction cycles for the remaining residues.

The use of various N-protecting groups, various coupling reagents, e.g., dicyclohexylcarbodiimide or carbonyldiimidazole, various active esters, e.g., esters of N-hydroxyphthalimide or N-hydroxysuccinimide, and the various cleavage reagents, to carry out reaction in solution, with subsequent isolation and purification of intermediates, is well known classical peptide methodology. Classical solution synthesis is described in detail in the treatise "Methoden der Organischen Chemie (Houben-Weyl): Synthese von Peptiden", E. Wunsch (editor) (1974) Georg Thieme Verlag, Stuttgart, W. Ger. Techniques of exclusively solid-phase synthesis are set forth in the textbook "Solid-Phase Peptide Synthesis", Stewart & Young, Pierce Chemical Co., Rockford, Ill., 1984, and are exemplified by the disclosure of U.S. Pat. No. 4,105,603. The fragment condensation method of synthesis is exemplified in U.S. Pat. No. 3,972,859. Other available syntheses are exemplified by U.S. Pat. Nos. 3,842,067 and 3,862,925.

Peptides are preferably prepared using the Merrifield solid phase synthesis, although other equivalent chemical syntheses known in the art can also be used as previously mentioned. Such solid-phase synthesis is commenced from the C-terminus of the peptide by coupling a protected *alpha*-amino acid to a suitable resin. Such a starting material can be prepared by attaching an *alpha*-amino-protected amino acid by an ester linkage to a chloromethylated resin or a hydroxymethyl resin, or by an amide bond to a benzhydrylamine (BHA) resin or paramethylbenzhydrylamine (MBHA) resin. The preparation of the hydroxymethyl resin is described by Bodansky et al., Chem. Ind. (London) 38, 1597-98 (1966). Chloromethylated resins are commercially available from Bio Rad Laboratories, Richmond, Calif. and from Lab. Systems, Inc. The preparation of such a resin is described by Stewart et al., "Solid Phase Peptide Synthesis", supra.

The C-terminal amino acid, protected by Boc and by a side-chain protecting group, if appropriate, can be first coupled to a chloromethylated resin according to the procedure set forth in Chemistry Letters, K. Horiki et al. 165-168 (1978), using KF in DMF

at about 60°C. for 24 hours with stirring, when a peptide having free acid at the C-terminus is to be synthesized.

5 Conditions for removal of specific *alpha*-amino protecting groups may be used as described in Schroder & Lubke, "The Peptides", 1 pp 72-75, Academic Press (1965).

10 Activating reagents and their use in peptide coupling are described by Schroder & Lubke *supra*, in Chapter III and by Kapoor, J. Phar. Sci., 59, pp 1-27 (1970).

15 The success of the coupling reaction at each stage of the synthesis, if performed manually, is preferably monitored by the ninhydrin reaction, as described by E. Kaiser et al., Anal. Biochem. 34, 595 (1970). The coupling reactions can be performed automatically, as on a Beckman 990 automatic synthesizer, using a program such as that reported in Rivier et al. Biopolymers, 1978, 17, pp 1927-1938.

20 After completing the growing peptide chains, the protected peptide resin is treated with liquid hydrofluoric acid to deblock and release the peptides from the support. For preparing an amidated peptide, the resin support used in the synthesis is selected to supply a C-terminal amide, after peptide cleavage from the resin.  
25 After removal of the hydrogen fluoride, the peptide is extracted into 1M acetic acid solution and lyophilized.

0 The peptide can be isolated by an initial separation by gel filtration, to remove peptide dimers and higher molecular weight polymers, and also to remove undesired salts.

5 Test and tolerogenic peptide sequences need not correspond exactly to the amino acid sequence of the agent infecting the host from which the PBMCs to be tolerated are derived. It is well known that proteins from wild type isolates of infectious agents often contain differences relative to the sequences of reference peptides of that

agent. However, use of peptides synthesised according to reference sequences will typically provide the desired tolerogenic effects.

5 In some circumstances, it may be desirable and feasible to use a test or tolerogenic sequence not from the agent infecting the host, but from a related agent, as long as the agents are sufficiently closely related for immunological cross-reactivity to occur, such that the desired tolerance is induced.

10 Alternatively, it may be desirable deliberately to introduce sequence mutations relative to either a wild type isolate or reference isolate. For example, without wishing to be bound by any particular theory, it is believed that the test/tolerogenic sequences may exert their effects by being presented to T cells  
15 with a Tr1 phenotype (3) by antigen presenting cells. Therefore it may be desirable to introduce mutations into a tolerogenic peptide from a given infectious agent in order to enable it to bind to a broader range of MHC molecules, and thus be used to tolerise a larger proportion of a population towards target antigens.

20 Therefore test or tolerogenic peptides may be used which differ from known or wild type sequences for the corresponding region of the infectious agent protein, as long as they retain sufficient tolerogenic capability. This can readily be determined by use of  
25 the methods of the present invention.

Variant peptides can be produced by a mixture of conservative variation, i.e. substitution of one hydrophobic residue such as isoleucine, valine, leucine or methionine for another, or the  
30 substitution of one polar residue for another, such as arginine for lysine, glutamic for aspartic acid, or glutamine for asparagine. As is well known to those skilled in the art, altering the primary structure of a polypeptide by a conservative substitution may not significantly alter the activity of that peptide because the  
35 side-chain of the amino acid which is inserted into the sequence may be able to form similar bonds and contacts as the side chain of the amino acid which has been substituted out. This is so even when

the substitution is in a region which is critical in determining peptide conformation. Also included are variants having non-conservative substitutions. As is well known to those skilled in the art, substitutions to regions of a peptide which are not critical in determining its conformation may not greatly affect its activity because they do not greatly alter the peptide's three dimensional structure, and so may not affect the desired activity, e.g. MHC binding. In regions which are critical in determining the peptides conformation or activity such changes may confer advantageous properties on the polypeptide. Indeed, changes such as those described above may confer slightly advantageous properties on the peptide e.g. altered stability or specificity.

Generally variant peptides may be extended at the N- or C-termini, and the C-terminus may be amidated or have a free acid form.

A peptide which is an amino acid sequence variant will generally share at least about 50%, 60%, 70%, 80%, 90% or more sequence identity with a wild type or reference sequence from the relevant infectious agent. In this connection, "sequence identity" means strict amino acid identity between the sequences being compared.

Once an amino acid substitution or other modification is made as described above, the peptide is screened for the requisite tolerogenic activity, as described above.

As described above, compositions of the present invention may comprise, in addition to the tolerogenic peptide sequences and optionally target antigens, a pharmaceutically acceptable excipient, carrier, buffer, stabiliser or other materials well known to those skilled in the art. Such materials should be non-toxic and should not interfere with the efficacy of the active ingredient. The precise nature of the carrier or other material may depend on the route of administration, e.g. oral, intravenous, cutaneous or subcutaneous, nasal, intramuscular, intraperitoneal

Pharmaceutical compositions for oral administration may be in tablet, capsule, powder or liquid form. A tablet may include a solid carrier such as gelatin or an adjuvant. Liquid pharmaceutical compositions generally include a liquid carrier such as water, petroleum, animal or vegetable oils, mineral oil or synthetic oil. Physiological saline solution, dextrose or other saccharide solution or glycols such as ethylene glycol, propylene glycol or polyethylene glycol may be included as required.

As the compositions of the present invention comprise peptides as active agents, they will typically be delivered by other routes, e.g. by intravenous, cutaneous or subcutaneous injection, or injection at the site of affliction, when the active ingredient will be in the form of a parenterally acceptable aqueous solution which is pyrogen-free and has suitable pH, isotonicity and stability. Those of relevant skill in the art are well able to prepare suitable solutions using, for example, isotonic vehicles such as Sodium Chloride Injection, Ringer's Injection, Lactated Ringer's Injection. Preservatives, stabilisers, buffers, antioxidants and/or other additives may be included, as required.

For delayed release, the active agents, e.g. tolerogenic peptide sequences and target antigens, may be included in a pharmaceutical composition formulated for slow release, such as in microcapsules formed from biocompatible polymers or in liposomal carrier systems according to methods known in the art.

For continuous release of peptides, the peptides may be covalently conjugated to a water soluble polymer, such as a polylactide or biodegradable hydrogel derived from an amphipathic block copolymer, as described in U.S. Pat. No. 5,320,840. Collagen-based matrix implants, such as described in U.S. Pat. No. 5,024,841, are also useful for sustained delivery of peptide therapeutics. Also useful, particularly for subdermal slow-release delivery, is a composition that includes a biodegradable polymer that is self-curing and that forms an implant in situ, after delivery in liquid form. Such a

composition is described, for example in U.S. Pat. No. 5,278,202.

Thus in a further aspect, the present invention provides a pharmaceutical composition comprising a tolerogenic peptide-  
5 encoding nucleic acid molecule and its use in methods of therapy or diagnosis. The composition may further comprise a target antigen-encoding nucleic acid molecule, which may be contiguous with the tolerogenic peptide-encoding nucleic acid molecule.

10 In a further aspect, the present invention provides a pharmaceutical composition comprising one or more tolerogenic peptide sequences as defined above and its use in methods of therapy or diagnosis. The composition may further comprise one or  
15 more target antigens.

In further aspects, the present invention provides the above described tolerogenic peptide sequences and encoding nucleic acid molecules for use in the preparation of medicaments for therapy.

20 Peptides may preferably be administered by transdermal iontophoresis. One particularly useful means for delivering compounds is transdermal delivery. This form of delivery can be effected according to methods known in the art. Generally,  
transdermal delivery involves the use of a transdermal "patch"  
5 which allows for slow delivery of compound to a selected skin region. Such patches are generally used to provide systemic delivery of compound. Examples of transdermal patch delivery systems are provided by U.S. Pat. No. 4,655,766 (fluid-imbibing osmotically driven system), and U.S. Pat. No. 5,004,610 (rate  
0 controlled transdermal delivery system).

For transdermal delivery of peptides, transdermal delivery may preferably be carried out using iontophoretic methods, such as described in U.S. Pat. No. 5,032,109 (electrolytic transdermal  
5 delivery system), and in U.S. Pat. No. 5,314,502 (electrically  
transdermal delivery system).

For transdermal delivery, it may be desirable to include permeation enhancing substances, such as fat soluble substances (e.g., aliphatic carboxylic acids, aliphatic alcohols), or water soluble substances (e.g., alkane polyols such as ethylene glycol, 1,3-propanediol, glycerol, propylene glycol, and the like). In addition, as described in U.S. Pat. No. 5,362,497, a "super water-absorbent resin" may be added to transdermal formulations to further enhance transdermal delivery. Examples of such resins include, but are not limited to, polyacrylates, saponified vinyl acetate-acrylic acid ester copolymers, cross-linked polyvinyl alcohol-maleic anhydride copolymers, saponified polyacrylonitrile graft polymers, starch acrylic acid graft polymers, and the like. Such formulations may be provided as occluded dressings to the region of interest, or may be provided in one or more of the transdermal patch configurations described above.

In other treatment methods, the modulators may be given orally or by nasal insufflation, according to methods known in the art. For administration of peptides, it may be desirable to incorporate such peptides into microcapsules suitable for oral or nasal delivery, according to methods known in the art.

Administration is preferably in a "prophylactically effective amount" or a "therapeutically effective amount" (as the case may be, although prophylaxis may be considered therapy), this being sufficient to show benefit to the individual. The actual amount administered, and rate and time-course of administration, will depend on the nature and severity of what is being treated. Prescription of treatment, e.g. decisions on dosage etc, is within the responsibility of general practitioners and other medical doctors, and typically takes account of the disorder to be treated, the condition of the individual patient, the site of delivery, the method of administration and other factors known to practitioners. Examples of the techniques and protocols mentioned above can be found in Remington's Pharmaceutical Sciences, 16th edition, Osol, A. (ed), 1980.



Alternatively, targeting therapies may be used to deliver the active agent more specifically to certain types of cell, by the use of targeting systems such as antibody or cell specific ligands. Targeting may be desirable for a variety of reasons; for example if the agent is unacceptably toxic, or if it would otherwise require too high a dosage, or if it would not otherwise be able to enter the target cells.

Instead of administering these agents directly, they could be produced in the target cells by expression from an encoding gene introduced into the cells, e.g. in a viral vector. The vector could be targeted to the specific cells to be treated, or it could contain regulatory elements which are switched on more or less selectively by the target cells.

A composition may be administered alone or in combination with other treatments, either simultaneously or sequentially dependent upon the condition to be treated.

Specific embodiments of the invention will now be described in more detail, by way of example and not limitation, by reference to the accompanying drawings.

#### Brief description of the drawings

Figure 1 shows cytokine and proliferative responses of PBMC from healthy EBV seropositive and seronegative donors (19) to purified LMP1 (20). Representative results are shown from two EBV seropositive donors (n=10) and two seronegative donors.

Figure 2 shows cytokine and proliferative responses of PBMC from EBV seropositive donors to a panel of LMP1 peptides. Representative results obtained from one donor (n=20) are shown for cytokine ELISAs (IL-10, IL-4, gamma-IFN) and a proliferation assay. The broken line on each chart shows the minimum level considered to be a positive response (0.0).

Figure 3 shows a summary of the percentage of EBV seropositive donors (n=20) whose PBMC responded to each LMP1 peptide with cytokine secretion (IL-10, IL-4, gamma-IFN) or proliferation. The results were demonstrated to be reproducible by retesting all of the 18 available donors.

Figure 4 shows flow cytometric analysis (23) of the phenotype of IL-10 synthesizing cells. After gating on CD3<sup>+</sup> cells, cultured cells from two EBV seropositive donors (A+B and C+D) were analyzed for expression of CD4 and IL-10, with the % of double positive cells shown in the upper right quadrant of each panel. A+C were obtained from unstimulated cultures and B+D from cells stimulated with peptides P14 (aa 66-85) and P8 (aa 36-55) respectively (shown to induce IL-10 in these donors).

Figure 5 shows proliferative and  $\gamma$ -IFN responses by PBMC from EBV seropositive, but not seronegative, donors against a mitogen (Con A), a recall antigen (PPD) and a primary antigen (KLH) (33) in the presence and absence of LMP1. Representative results are shown from two EBV seropositive donors (n=10) and two seronegative donors.

Figure 6 shows that IL-10 inducing LMP1 peptides inhibit proliferative responses by PBMC from EBV seropositive donors against recall antigen (PPD). The white bars show the proliferative and gamma-IFN responses obtained when PBMC from three EBV seropositive donors were stimulated with PPD, either alone, or together with IL-10 inducing LMP1 peptides (P4,7,23,35 for Donor 1, P4 and 22 for Donor 2, and P4,18 and 31 for Donor 3), or control gamma-IFN inducing LMP1 peptides (P28 for Donor 1, P56 for Donor 2 and P33 for Donor 3). The black bars show the effects of adding a neutralizing anti-IL-10 antibody to duplicate cultures at 0.5 $\mu$ g/ml.

Figures 7 to 10 illustrate the specificity and persistence of LMP-1-induced tolerance. In each Figure, panel (a) shows proliferative,  $\gamma$ -IFN and IL-10 responses obtained when PBMC from a given donor were first stimulated in culture with the mitogen Con

A, the recall antigen PPD, or the primary antigen KLH, alone or in combination with purified LMP1. In Figures 7a and 9a, stimuli were also administered in combination with an LMP-1-derived peptide. Cells were rested for seven days, washed to remove the antigens, and added to fresh irradiated autologous PBMC as a source of antigen presenting cells (33). Panel (b) shows the results of restimulating the control cells with each of the three stimuli. Panel (c) shows the results of restimulating the cells originally stimulated in the presence of LMP-1. Panel (d), shown only for Figures 7 and 9, shows the results of restimulating cells originally stimulated in the presence of LMP-1-derived peptide. Results are shown for three EBV-seropositive donors (Figures 7, 8 and 9) and one seronegative donor (Figure 10).

### Examples

**LMP1 induces high levels of IL-10 secretion by PBMC from EBV seropositive but not seronegative donors.**

PBMC from ten EBV seropositive donors were tested for the ability to respond to purified LMP1 with either Th cytokine secretion or proliferation. In all seropositive donors, IL-10 was the predominant cytokine measured, with no significant proliferative, gamma-IFN or IL-4 responses. Figure 1 shows representative results obtained from two seropositive donors. To confirm that the observed responses resulted from previous EBV infection, PBMC from two EBV seronegative donors were tested for responsiveness to the purified LMP1. It can also be seen from Figure 1 that, in these donors, the LMP1 failed to elicit either IL-10 secretion, or significant proliferative and  $\gamma$ -IFN responses. The results in both donor groups are specific to LMP1, since the T-cell mitogen concanavalin A (Con A) and the control recall antigen *Mycobacterium tuberculosis* purified protein derivative (PPD) induced responses dominated by proliferation and  $\gamma$ -IFN production, regardless of EBV serological status.

FIGURE 1. EBV seropositive donors: PBMC stimulated with LMP1, Con A, or PPD. IL-10 secretion is high in LMP1-stimulated cells, but low in Con A- and PPD-stimulated cells. EBV seronegative donors: PBMC stimulated with LMP1, Con A, or PPD. IL-10 secretion is low in all three conditions.

To further characterize the immune response to LMP1, epitopes that induced IL-10 secretion were mapped by screening PBMC from 20 EBV seropositive healthy donors with a panel of synthetic 20-mer peptides spanning the entire sequence of LMP1. Representative results obtained from one donor (Figure 2) demonstrate that multiple LMP1 peptides induced secretion of high concentrations of IL-10. In contrast, only three peptides induced proliferation, five peptides  $\gamma$ -IFN, two peptides IL-4, and all the latter responses were weak. Similar patterns of responsiveness were found in a total of 20 seropositive donors (summarized in Figure 3). Strikingly, certain peptides commonly elicited IL-10 responses in different donors ( $p=2.2 \cdot 10^{-7}$ , Poisson heterogeneity test), with, for example, peptide 4 (aa 16-35) inducing IL-10 in 80% of the individuals. Furthermore, these dominant IL-10 inducing peptides are clustered within the N terminal half of the protein that is rich in binding motifs for many MHC class II molecules (<http://imtech.res.in/raghava/propred/index.html>; <http://www.csd.abdn.ac.uk/~gjlk/MHC-Thread>) (Sturniolo et al. (1999); Singh and Raghava (2001)).

PBMC from four EBV seronegative donors were also screened with the panel of LMP1 peptides. Reactivity was rare in this group, with totals of only nine IL-10, one  $\gamma$ -IFN, one proliferative and no IL-4 responses. Moreover, all these responses were relatively weak (data not shown).

**Cells responding to LMP1 and LMP1 peptides with IL-10 secretion are CD3<sup>+</sup>CD4<sup>+</sup>.**

The phenotype of the cells responsible for the IL-10 production was determined by flow cytometry, comparing peptide stimulated and unstimulated cultures from four seropositive individuals. Most IL-10 producing cells bore the CD3 marker for T-cells (mean=83.9%, SD=9.4%) and of these the majority were of the CD4<sup>+</sup> helper phenotype (mean=90.6%, SD=8.9%) (Figure 4). Similarly, activated cells, as judged by expression of CD69 and CD71, in the rare cultures proliferating in response to peptides were also CD4<sup>+</sup> (data not shown).

LMP1 and LMP1 peptides suppress proliferative and gamma-IFN responses by stimulating IL-10 secreting Tr1 cells.

There is evidence that CD4<sup>+</sup> T-cells biased towards IL-10 secretion, termed Tr1 cells, play an important role in immunoregulation (Groux et al., 1997; Levings and Roncarolo, 2000) and have been shown to inhibit inflammatory responses (Groux et al., 1997; Roncarolo and Levings, 2000). We concluded that the responses to LMP1 and the peptide panel were predominantly mediated by Tr1 cells and sought to confirm that they were capable of mediating suppression. In all ten seropositive donors tested, the addition of LMP1 strongly inhibited proliferative and gamma-IFN responses to the T-cell mitogen Con A, the recall antigen PPD and the primary antigen keyhole limpet hemocyanin (KLH) by 56-99% (Figure 5). The IL-10 responses to LMP1, and the associated inhibition, were dependent on the donor having been infected with EBV, since no such effects were seen when PBMC were obtained from two control seronegative volunteers (Figure 5). Results similar to those obtained with purified LMP1 protein were found when IL-10 inducing LMP1 peptides were added to PBMC cultures from five seropositive donors, with suppression of proliferative and gamma-IFN responses to PPD by 41-99% (Figure 6). In parallel experiments, the peptides also inhibited proliferative and gamma-IFN responses to the mitogen Con A or primary antigen KLH (results not shown). Figure 6 also demonstrates that the inhibitory effect is dependent on IL-10, since, when LMP1 derived peptides that did not elicit this cytokine were added to PPD-stimulated cultures, no suppression was seen. Furthermore, in cultures treated with anti-IL-10 antibody, the LMP1 peptide mediated suppression was reversed by up to 71% (Figure 6).

*In vitro* LMP1 mediated suppression is antigen specific and persistent.

PBMCs from three seropositive donors and one seronegative donor were first stimulated in culture with the mitogen Con A, the recall antigen PPD, or the primary antigen KLH, alone or in combination with purified LMP1 and in two cases in combination with an IL-10

washed to remove the antigens, and added to fresh irradiated autologous PBMC as a source of antigen presenting cells (Plebanski et al., 1992). Each group of cells was then restimulated with each stimulus. Results are shown in Figures 7 to 10.

5 Cells from seropositive donors (Figures 7, 8 and 9) which were initially exposed to a stimulus in combination with LMP-1 produced significant IL-10 and low IFN-gamma and proliferative responses. When re-stimulated with the same stimulus, in the absence of LMP-1, these cells still failed to proliferate or express IFN-gamma.  
 10 However, they retained the capacity to proliferate and produce IL-10 against other stimuli, showing that the cells had been specifically tolerised to the stimulus originally administered in combination with LMP-1. Similar results were obtained for two of the seropositive donors with LMP-1-derived peptides (peptides 4 and  
 15 18, shown in Figures 7d and 9d respectively).

Cells from the seronegative donor (Figure 10) responded with typical IFN-gamma and proliferative responses to all stimuli regardless of the presence of LMP-1, showing that the induction of tolerance is not an inherent property of the protein, but relies on  
 20 prior exposure of the cells to EBV.

Thus, it can be hypothesised that the Tr1 response to LMP1 deviates T-cells recognizing a bystander antigen to adopt an anergic, IL-10 secreting phenotype. Such induction of anergy specific for other viral antigens that are co-expressed with LMP1 may be important in  
 25 the maintenance of EBV latency.

#### Discussion of Examples

Here a novel mechanism is postulated by which Epstein-Barr virus (EBV), rather than avoiding detection, instead subverts the immune  
 30 response by stimulating regulatory CD4<sup>+</sup> T-cells that secrete the inhibitory cytokine interleukin-10 (IL-10). Such regulatory T-cells are well recognized (1-3) but not known to have a role in viral persistence (4-6). Human peripheral blood mononuclear cells (PBMC) from all EBV seropositive, but not seronegative, donors  
 35 responded to both purified latent membrane protein 1 (LMP1) and the

corresponding immunodominant peptides with high levels of IL-10 secretion by CD4<sup>+</sup> T-cells. These IL-10 responses, characteristic of T regulatory 1 (Tr1) cells, coincided with inhibition of T-cell proliferation and  $\gamma$ -interferon ( $\gamma$ -IFN) secretion induced by both mitogen and recall antigen. The ability of this viral antigen to deviate the immune response towards tolerance is likely to be important in maintaining latency and EBV associated tumors.

This study was prompted by the lack of a protective immune response against LMP1 in individuals with latent EBV infection. The main conclusion from our data is that LMP1 is recognized by the immune system, but that this response is dominated by the induction of high levels of IL-10 secretion by cells with a Tr1 phenotype (Levings and Roncarolo, 2000). Furthermore, this IL-10 response was able to suppress both proliferative and  $\gamma$ -IFN responses against other antigens and polyclonal stimuli, and therefore would be expected to prevent the development of protective Th1 and cytotoxic immunity against LMP1 (Fiorentino et al., 1991). Indeed, such IL-10 secretion is also likely to anergise Th1 responses to other EBV proteins co-expressed both in latent infection and associated tumors. Our demonstration of Tr1 activation provides a mechanism for the previously reported observation that recombinant LMP1 inhibits immune functions including mitogen, antigen and CD3/CD28 stimulated T-cell activation; natural killer cell cytotoxicity; and antigen-induced gamma-IFN secretion (Dukers et al., 2000). One peptide from LMP1, included within the sequence of peptide 7 (aa 31-50) from our panel, was reported (Dukers et al., 2000) to replicate these inhibitory properties, and is identified here as one of many effective Tr1 IL-10 inducers.

A possible explanation for the propensity of LMP1 to elicit IL-10 production by Tr1 cells is that the establishment and maintenance of a suppressive response to LMP1 results from IL-10 'conditioning'. Activation of CD4<sup>+</sup> T-cells in the presence of IL-10 leads to the generation of Tr1 cells (Groux et al., 1997). EBV

lytic cycle infection (Hayes et al., 1999). Thus, we propose that, during the development of the immune response to EBV, the presence of vIL-10 deviates the differentiation of LMP1 specific Th cells to favor IL-10 secreting Tr1 cells, a bias which then becomes self-perpetuating. The finding that the regulatory response to LMP1 is limited to EBV seropositive, but not seronegative, donors strongly supports this second explanation.

Numerous methods used by pathogens to avoid clearance by the immune system have been described, dependent on escape, resistance or counterattack (Xu et al., 2001). The induction of a Tr1 response to EBV LMP1 represents a further mechanism of immune evasion. A similar regulatory mechanism that subverts, rather than avoids, immune detection may well be exploited by other pathogens with the ability to maintain chronic infections. This is especially likely for those viruses, such as cytomegalovirus, that also encode a homologue of IL-10 with potent immunosuppressive effects (Spencer et al., 2002), which may also induce a regulatory Tr1 type immune response. The design of strategies to overcome such Tr1 responses should provide an innovative approach to the development of vaccines to prevent or treat EBV associated tumors. Conversely, it may be possible to exploit therapeutically such specific induction of bystander anergy to inhibit pathogenic responses in immune-mediated diseases.

## Experimental Procedures

### **Donors**

Blood samples were obtained by venepuncture from a group of healthy volunteers. The donors were classified as EBV seropositive or seronegative by an ELISA for serum anti-EBNA1 IgG, with negative results confirmed by immunofluorescence staining for IgG and IgM anti-viral capsid antibody.

---

### **Antigens and Mitogen**



LMP1 was immunopurified from lysed EBV transformed B cells using the anti-LMP1 antibody CS1-4 (Novocastra Laboratories) conjugated to anti-mouse IgG<sub>1</sub> coated magnetic beads (Biomag, PerSeptive Biosystems).

A panel of 76 20-mer peptides, with 15 amino acid overlaps, was synthesised (Department of Biochemistry, University of Birmingham, UK or University of Bristol, UK), spanning the entire length of the 63kD EBV LMP1, as determined from the prototype B-cell-derived gene (B95.8) sequence (Hayes et al., 1999). All peptides were used to stimulate cultures at 15µg/ml, although, as in previous mapping studies (Stott et al., 2000), responses were similar over a wide range of concentrations (4-50µg/ml).

Peptide sequences are as follows:

P1	MEHDLERGPPGPRRPPRGPP	P39	HSDEHHHDDSLPHPQQATDD
P2	ERGPPGPRRPPRGPPPLSSSL	P40	HHDDSLPHPQQATDDSGHES
P3	GPRRPPRGPPPLSSSLGLALL	P41	LPHPQQATDDSGHESDSNSN
P4	PRGPPPLSSSLGLALLLLLLLA	P42	QATDDSGHESDSNSNEGRHH
P5	LSSSLGLALLLLLLLALLFWL	P43	SGHESDSNSNEGRHHLLVSG
P6	GLALLLLLLLALLFWLYIVMS	P44	DSNSNEGRHHLLVSGAGDGP
P7	LLLLALLFWLYIVMSDWTGG	P45	EGRHHLLVSGAGDGPPLCSQ
P8	LLFWLYIVMSDWTGGALLVL	P46	LLVSGAGDGPPLCSQNLGAP
P9	YIVMSDWTGGALLVLYSFAL	P47	AGDGPPLCSQNLGAPGGGPD
P10	DWTGGALLVLYSFALMLIII	P48	PLCSQNLGAPGGGPDNGPQD
P11	ALLVLYSFALMLIIIIILIIIF	P49	NLGAPGGGPDNGPQDPDNTD
P12	YSFALMLIIIIILIIIFIFRRD	P50	GGGPDNGPQDPDNTDDNGPQ
P13	MLIIIIILIIIFIFRRDLLCPL	P51	NGPQDPDNTDDNGPQDPDNT
P14	ILIIIFIFRRDLLCPLGALCI	P52	PDNTDDNGPQDPDNTDDNGP
P15	IFRRDLLCPLGALCILLMI	P53	DNGPQDPDNTDDNGPHDPLP
P16	LLCPLGALCILLMITLLLI	P54	DPDNTDDNGPHDPLPQDPDN
P17	GALCILLMITLLLIALWNL	P55	DDNGPHDPLPQDPDNTDDNG
P18	LLLMITLLLIALWNLHGQAL	P56	HDPLPQDPDNTDDNGPQDPD
P19	TLLLIALWNLHGQALFLGIV	P57	QDPDNTDDNGPQDPDNTDDN
P20	ALWNLHGQALFLGIVLFIFG	P58	TDDNGPQDPDNTDDNGPHDP
P21	HGQALFLGIVLFIFGCLLVL	P59	PQDPDNTDDNGPHDPLPHSP
P22	FLGIVLFIFGCLLVLGIIWY	P60	NTDDNGPHDPLPHSPSDSAG
P23	LFIFGCLLVLGIIWYLLLEML	P61	GPHDPLPHSPSDSAGNDGGP
P24	CLLVLGIIWYLLLEMLWRLGA	P62	LPHSPSDSAGNDGGPPQLTE
P25	GIWIYLLLEMLWRLGATIWQL	P63	SSGSGGDDDDPHGVPQLSYYD
P26	LLEMLWRLGATIWQLLAFFL	P64	SDSAGNDGGPPQLTEEVENK
P27	WRLGATIWQLLAFFLAFFLD	P65	NDGGPPQLTEEVENKGGDQG
P28	TIWQLLAFFLAFFLDLILLI	P66	PQLTEEVENKGGDQGPPLMT
P29	TAFTTATFLDILLIILALIL	P67	EVENKGGDQGPPLMTDGGGG
P30	TAFTTATFLDILLIILALIL		
P31	TAFTTATFLDILLIILALIL		
P32	TAFTTATFLDILLIILALIL		
P33	TAFTTATFLDILLIILALIL		
P34	TAFTTATFLDILLIILALIL		
P35	TAFTTATFLDILLIILALIL		
P36	TAFTTATFLDILLIILALIL		
P37	TAFTTATFLDILLIILALIL		
P38	TAFTTATFLDILLIILALIL		

P33 QQNWWTLLVDLLWLLLFLAI  
 P34 TLLVDLLWLLLFLAILIWMY  
 P35 LLWLLLFLAILIWMYYHGQR  
 P36 LFLAILIWMYYHGQRHSDEH  
 P37 LIWMYYHGQRHSDEHHHDDS  
 P38 YHGQRHSDEHHHDDSLPHPQ

P71 HSHDSGHGGGDPHLPTLLLG  
 P72 GHGGGDPHLPTLLLGSSGSG  
 P73 DPHLPTLLLGSSGSGGDDDD  
 P74 TLLGSSGSGGDDDDDPHGPV  
 P75 LFLAILIWMYYHGQRHSDEH

A similar panel of peptides was prepared from LMP2 having sequences as follows:

P1' MGSLEMVPMGAGPPSPGGDP  
 P2' MVPMGAGPPSPGGDPDGYDG  
 P3' AGPPSPGGDPDGYDGGNNSQ  
 P4' PGGDPDGYDGGNNSQYPSAS  
 P5' DGYDGGMNSQYPSASGSSGN  
 P6' GNSQYPSASGSSGNTPTPP  
 P7' YPSASGSSGNTPTPPMDEER  
 P8' GSSGNTPTPPNDEERESNEE  
 P9' TPTPPNDEERESNEEPPPPY  
 P10' NDEERESNEEPPPPYEDPYW  
 P11' RESNEEPPPPYEDYWGNGD  
 P12' EPPPPYEDPYWGNGDRHSDY  
 P13' YEDPYWGNGDRHSDYQPLGT  
 P14' WGNGDRHSDYQPLGTQDQSL  
 P15' RHSDYQPLGTQDQSLYLGLQ  
 P16' QPLGTQDQSLYLGLQHDGND  
 P17' QDQSLYLGLQHDGNDGLPPP  
 P18' LGLQHDGNDGLPPPYSPRD  
 P19' DGNDGLPPPYSPRDDSSQH  
 P20' LPPPPYSPRDDSSQHIYEEA  
 P21' PPYSPRDDSSQHIYEEAGRG  
 P22' RDDSSQHIYEEAGRGSMNPV  
 P23' QHIYEEAGRGSMNPVCLPVI  
 P24' EAGRGSMMPVCLPVIAPYL  
 P25' SMNPVCLPVIAPYLFWLAA  
 P26' CLPVIAPYLFWLAAIAASC  
 P27' VAPYLFWLAAIAASCFTASV  
 P28' FWLAAIAASCFTASVSTVVT  
 P29' IAASCFTASVSTVVTATGLA  
 P30' FTASVSTVVTATGLALSLLL  
 P31' STVVTATGLALSLLLLAAVA  
 P32' ATGLALSLLLLAAVASSYAA  
 P33' LSLLLLAAVASSYAAAQRKL  
 P34' LAAVASSYAAAQRKLLTPVT  
 P35' SSYAAAQRKLLTPVTVLTAV  
 P36' AQRKLLTPVTVLTAVVTFFA  
 P37' LTPVTVLTAVVTFFAICLTW  
 P38' VLTAVVTFFAICLTWRIEDP  
 P39' VTFFAICLTWRIEDPPFNSL  
 P40' ICLTWRIEDPPFNSLLFALL  
 P41' RIEDPPFNSLLFALLAAAGG  
 P42' PFNSLLFALLAAAGGLQGIY  
 P43' LFALLAAAGGLQGIYVLVML  
 P44' AAAGGLQGIYVLVMLVLLIL

P49' WRRITVCGGIMFLACVLVLI  
 P50' VCGGIMFLACVLVLIVDAVL  
 P51' MFLACVLVLIVDAVLQLSPL  
 P52' VLVLIIVDAVLQLSPLLGAVT  
 P53' VDAVLQLSPLLGAVTVVSMT  
 P54' QLSPLLGAVTVVSMTLLLLA  
 P55' LGAVTVVSMTLLLLAFVLWL  
 P56' VVSMTLLLLAFVLWLSSPGG  
 P57' LLLLLAFVLWLSSPGGLGTLG  
 P58' FVLWLSSPGGLGTLGAALLT  
 P59' SSPGGLGTLGAALLTLAAAL  
 P60' LGTLGAALLTLAAALALLAS  
 P61' AALLTLAAALALLASLILGT  
 P62' LAAALALLASLILGTINLTT  
 P63' ALLASLILGTINLTTMFLLM  
 P64' LILGTINLTTMFLLMLLWTL  
 P65' LNLTMTFLMLLWTLVLLI  
 P66' MFLMLLWTLVLLIICSSCS  
 P67' LLWTLVLLIICSSCSCPLS  
 P68' VLLIICSSCSCPLSKILLA  
 P69' CSSCSCPLSKILLARFLY  
 P70' SCPLSKILLARFLYALALL  
 P71' KILLARFLYALALLLLASA  
 P72' RLFLYALALLLLASALIAGG  
 P73' ALALLLLASALIAGGSILQT  
 P74' LLASALIAGGSILQTNFKSL  
 P75' LIAGGSILQTNFKSLSSSTEF  
 P76' SILQTNFKSLSSSTEFIPNLF  
 P77' NFKSLSSSTEFIPNLFMLLL  
 P78' SSTEIPNLFMLLLIVAGI  
 P79' IPNLFMLLLIVAGILFILAI  
 P80' CMLLLIVAGILFILAILTEW  
 P81' IVAGILFILAILTEWGSGNR  
 P82' LFILAILTEWGSGNRTYGPV  
 P83' ILTEWGSGNRTYGPVFMCLG  
 P84' GSGNRTYGPVFMCLGGLLTM  
 P85' FMCLGGLLTMVAGAVWLTVM  
 P86' GLLTMVAGAVWLTVMSTLL  
 P87' VAGAVWLTVMSTLLSAWIL  
 P88' WLTVMSTLLSAWILTAGFL  
 P89' SNTLLSAWILTAGFLIFLIG  
 P90' SAWILTAGFLIFLIGFALFG  
 P91' TAGFLIFLIGFALFGVIRCC  
 P92' IFLIGFALFGVIRCCRYCCY

P45'	LQGIYVLVMLVLLILAYRRR	P93'	FALFGVIRCCRYCCYYCLTL
P46'	VLVMLVLLILAYRRRWRLT	P94'	VIRCCRYCCYYCLTLESEER
P47'	VLLILAYRRRWRLTVCGGI	P95'	RYCCYYCLTLESEERPPTY
P48'	AYRRRWRLTVCGGIMFLAC	P96'	YYCLTLESEERPPTYRNTV

The control antigen mycobacterial PPD (Statens Seruminstitut), the T-cell mitogen Con A (Sigma), and the primary antigen keyhole limpet hemocyanin (KLH) (Calbiochem) were each used to stimulate cultures at 10µg/ml. PPD readily provokes recall T-cell responses *in vitro* (Plebanski et al., 1992), since most UK citizens have been immunised with Bacille Calmette-Guérin (BCG).

#### Cell Proliferation and Cytokine Assays

As described elsewhere (Stott et al., 2000), PBMC were separated from fresh blood samples by density gradient centrifugation and cultured in 1ml volumes at a concentration of  $1.25 \times 10^6$  cells/ml. Cellular proliferation was estimated from the incorporation of  $^3\text{H}$ -thymidine in triplicate 100µl aliquots taken from the wells five days after antigen stimulation, when recall T-cell responses are maximal (Plebanski et al., 1992). Proliferation results are presented as the mean counts per minute (CPM)  $\pm$  SD of the triplicate samples, or as stimulation index (SI), expressing the ratio of mean CPM in stimulated versus unstimulated control cultures. An  $\text{SI} > 3$  with  $\text{CPM} > 1000$  is interpreted as representing a significant positive response (Devereux et al., 2000). Production of the Th cytokines  $\gamma$ -IFN, IL-4 and IL-10 was assessed in duplicate 100µl aliquots taken five days after stimulation of the cultures, using a sensitive cellular ELISA (Devereux et al., 2000). Cytokine responses over twice the production in unstimulated cultures were considered positive (Devereux et al., 2000).

#### Characterisation of Responding Cells

The phenotypes of cultured cells that proliferate or secrete cytokine in response to antigen were determined by flow cytometry.

Aliquots of PBMC were taken from responding cultures and stained

isothiocyanate, with anti-CD25 phycoerythrin-cyanin 5.1 (all Beckman Coulter) in some experiments. Activated cells in proliferating cultures were identified using anti-CD71-phycoerythrin (PE) or anti-CD69-PE (both Beckman Coulter). Cells synthesising IL-10 were labelled by incubating with anti-IL-10-PE (Pharmingen) after inhibition of protein secretion with Brefeldin A (Sigma) and permeabilisation with Intraprep™ (Beckman Coulter). Stained cells were analysed using an EPICS XL cytometer (Beckman Coulter) and Expo v2 analysis software (Applied Cytometry Systems).

10 All documents cited herein are hereby incorporated by reference insofar as is required for the skilled person to carry out the invention.

## References

Apostolopoulos V, Plebanski M. (2000) The evolution of DNA vaccines. *Curr Opin Mol Ther* 2(4):441-7

5 Baer, R., Bankier, A.T., Biggin, M.D., Deininger, P.L., Farrell, P.J., Gibson, T.J., Hatfull, G., Hudson, G.S., Satchwell, S.C., Seguin, C., et al. (1984). DNA sequence and expression of the B95-8 Epstein-Barr virus genome. *Nature* 310, 207-11.

10 Blaschke V, Reich K, Blaschke S, Zipprich S, Neumann C. (2000) Rapid quantitation of proinflammatory and chemoattractant cytokine expression in small tissue samples and monocyte-derived dendritic cells: validation of a new real-time RT-PCR technology. *J Immunol Methods* 246(1-2):79-90.

15 Chapman, A.L.N., Rickinson A.B., Thomas W.A., Jarrett R.F., Crocker J., and Lee S.P. (2001). Epstein-Barr virus-specific cytotoxic T lymphocyte responses in the blood and tumor site of Hodgkin's disease patients: implications for a T-cell-based therapy. *Cancer Res.* 61, 6219-26.

20 Christensen, J.P., Bartholdy, C., Wodarz, D., and Thomsen, A.R. (2001). Depletion of CD4+ T cells precipitates immunopathology in immunodeficient mice infected with a noncytotoxic virus. *J. Immunol.* 166(5), 3384-91.

Demay F, Darche S, Kourilsky P, Delassus S. (1996) Quantitative PCR with colorimetric detection. *Res Immunol* 147(6):403-6.

25 Devereux, G., Hall, A.M., and Barker, R.N. (2000). Measurement of T-helper cytokines secreted by cord blood mononuclear cells in response to allergens. *J Immunol Methods.* 234, 13-22.

0 Dieckmann, D., Plottner, H., Berchtold, S., Berger, T., and Schuler, G. (2001). *Ex vivo* isolation and characterisation of CD4+CD25+ T cells with regulatory properties from human blood. *J. Exp. Med.* 193, 1303-1310.

Palmer, A. F. (2001) T helper 1 and T helper 2 cells: a review. *Immunol Rev.* 181, 1-11.

immunosuppressive effects of EBV-encoded latent membrane protein 1. *J Immunol.* 165, 663-670.

5 Eliopoulos, A.G., Dawson, C.W., Mosialos, G., Floettmann, J.E., Rowe, M., Armitage, R.J., Dawson, J., Zapata, J.M., Kerr, D.J., Wakelam, M.J., Reed, J.C., Kieff, E., and Young, L.S. (1996). CD40-induced growth inhibition in epithelial cells is mimicked by Epstein-Barr Virus-encoded LMP1: involvement of TRAF3 as a common mediator. *Oncogene.* 13, 2243-2254.

10 Eliopoulos, A.G., Stack, M., Dawson, C.W., Kaye, K.M., Hodgkin, L., Sihota, S., Rowe, M., and Young, L.S. (1997). Epstein-Barr virus-encoded LMP1 and CD40 mediate IL-6 production in epithelial cells via an NF-kappaB pathway involving TNF receptor-associated factors. *Oncogene.* 14, 2899-2916.

15 Erickson, A.L., Kimura, Y., Igarashi, S., Eichelberger, J., Houghton, M., Sidney, J., McKinney, D., Sette, A., Hughes, A.L., and Walker, C.M. (2001). The outcome of Hepatitis C virus infection is predicted by escape mutation in epitopes targeted by cytotoxic T lymphocytes. *Immunity* 15, 883-895.

20 Fiorentino, D.F., Zlotnik, A., Vieira, P., Mosmann, T.R., Howard, M., Moore, K.W., O'Garra, A. (1991) IL-10 acts on the antigen-presenting cell to inhibit cytokine production by Th1 cells. *J Immunol.* 146, 3444-3451.

25 Fries, K.L., Miller, W.E., and Raab-Traub, N. (1996). Epstein-Barr virus latent membrane protein 1 blocks p53-mediated apoptosis through the induction of the A20 gene. *J. Virol.* 70, 8653-8659.

Gorelik, L. and Flavell, R.A. (2000). Abrogation of TGF beta signalling in T cells leads to spontaneous T cell differentiation and autoimmune disease. *Immunity* 12, 171-181.

Gregory, C.D., Murray, R.J., Edwards, C.F., and Rickinson, A.B.

~~30~~ (1988) ~~Downregulation of cell adhesion molecules LFA-3 and ICAM-1~~ in Epstein-Barr virus-positive Burkitt's lymphoma underlies tumor cell escape from virus-specific T cell surveillance. *J. Exp. Med.* 167, 1811-1824.

Groux, H., O'Garra, A., Bigler, M., Rouleau, M., Antonenko, S., de Vries, J.E., and Roncarolo, M.G. (1997). A CD4+ T-cell subset inhibits antigen-specific T-cell responses and prevents colitis. *Nature*. 389 (6652), 737-742.

5 Haraguchi, S., Good, R.A., and Day, N.K. (1995) Immunosuppressive retroviral peptides: cAMP and cytokine patterns. *Immunol. Today* 16, 595.

Hayes, D.P., Brink, A.A., Vervoort, M.B., Middeldorp, J.M., Meijer, C.J., van den Brule, A.J (1999) Expression of Epstein-Barr virus  
10 (EBV) transcripts encoding homologues to important human proteins in diverse EBV associated diseases. *Mol Pathol*. 52, 97-103.

Horikawa, T., Yoshizaki, T., Sheen, T.S., Lee, S.Y., and Furukawa, M. (2000). Association of latent membrane protein 1 and matrix metalloproteinase 9 with metastasis in nasopharyngeal carcinoma.  
15 *Cancer*. 89, 715-23.

Huen, D.S., Henderson, S.A., Croom-Carter, D., and Rowe, M. (1995). The Epstein-Barr virus latent membrane protein-1 (LMP1) mediates activation of NF-kappa B and cell surface phenotype via two effector regions in its carboxy-terminal cytoplasmic domain.  
20 *Oncogene*. 10, 549-560.

Jonuleit, H., Schmitt, E., Stassen, M., Tuettenberg, A., Knop, J., and Enk, A. (2001). Identification and Functional Characterization of Human CD4+CD25+ T Cells with Regulatory Properties Isolated from Peripheral Blood. *J. Exp. Med*. 193, 1285-1294.

5 Kamel-Reid S, Dick JE. Engraftment of immune-deficient mice with human hematopoietic stem cells. *Science* 1988;242:1706-9.

Khanna, R., Burrows, S.R., Nicholls, J., and Poulsen, L.M. (1998). Identification of cytotoxic T cell epitopes within Epstein-Barr virus (EBV) oncogene latent membrane protein 1 (LMP1): evidence for  
0 HLA A2 supertype-restricted immune recognition of EBV-infected cells by LMP1-specific cytotoxic T lymphocytes. *Int J Cancer* 77, 1111-1118.

Kieff, E. (1996) Epstein-Barr virus and its replication. In Fields Virology, Fields B. N., Knipe, D. M. and Howley, P. M eds. (Lipincott-Raven Publishers, Philadelphia), pp2343-2396.

- 5 Kjellstrom S, Emneus J, Marko-Varga G. (2000) Flow immunochemical bio-recognition detection for the determination of interleukin-10 in cell samples. J Immunol Methods 246(1-2):119-30.
- 10 Kreft B, Singer GG, Diaz-Gallo C, Kelley VR. (1992) Detection of intracellular interleukin-10 by flow cytometry. J Immunol Methods 156(1):125-8
- Levings, M.K. and Roncarolo, M.G. (2000). T-regulatory 1 cells: a novel subset of CD4 T cells with immunoregulatory properties. J Allergy Clin Immunol. 106, S109-112.
- 15 Levings, M.K., Sangregorio, R., and Roncarolo, M.G. (2001). Human CD25+CD4+ T regulatory cells suppress naïve and memory T cell proliferation and can be expanded *in vitro* without loss of function. J. Exp. Med. 193, 1295-1302.
- 20 Levitskaya, J., Coram, M., Levitsky, V., Imreh, S., Steigerwald-Mullen, P.M., Klien, G., Kurilla, M.G., and Masucci, M.G. (1995). Inhibition of antigen processing by the internal repeat region of the Epstein-Barr virus nuclear antigen-1. Nature 375, 685-688.
- 25 Levitskaya, J., Sharipo, A., Leonchiks, A., Ciechanover, A., and Masucci, M.G. (1997). Inhibition of ubiquitin/proteasome- dependent protein degradation by the Gly-Ala repeat domain of the Epstein-Barr virus nuclear antigen 1. Proc. Natl. Acad. Sci. USA 94, 12616-12621.
- 30 Lorenzo, M.E., Ploegh, H.L., and Tirabassi, R.S. (2001). Viral immune evasion strategies and the underlying cell biology. Semin Immunol. 13, 1-9.
- MacDonald, AJ, Duffy, M, Brady, MT, McKiernan, S, Hall, W, Hegarty, J, Curry, M, and Mills KHG. CD4 T Helper Type 1 and Regulatory T Cells Induced against the Same Epitopes on the Core Protein in



Hepatitis C Virus-Infected Persons. The Journal of Infectious Diseases (2002) 185:720-7

McCune JM, Namikawa R, Kaneshima H, Shultz LD, Lieberman M, Weissman IL. The SCID-hu mouse: murine model for the analysis of human hematolymphoid differentiation and function. Science 1988;241:1632-9.

Mor G, Eliza M. (2001) Plasmid DNA vaccines. Immunology, tolerance, and autoimmunity. Mol Biotechnol 19(3):245-50

Mosialos, G., Birkenbach, M., Yalamanchili, R., VanArsdale, T., Ware, C., and Kieff, E. (1995). The Epstein-Barr virus transforming protein LMP1 engages signaling proteins for the tumor necrosis factor receptor family. Cell. 80, 389-399.

Mosier DE, Gulizia RJ, Baird SM, Wilson DB. Transfer of a functional human immune system to mice with severe combined immunodeficiency. Nature 1988 335:256-9

Mossman, T.R., and Coffman, R.L. (1989). Heterogeneity of cytokine secretion patterns and functions of helper T-cells. Adv. Immunol. 46, 111-147.

Mueller, Y.M., De Rosa, S.C., Hutton, J.A., Witek, J., Roederer, M., Altman, J.D., and Katsikis, P.D. (2001). Increased CD95/Fas-Induced apoptosis of HIV-Specific CD8<sup>+</sup> T cells. Immunity 15, 883-895.

Pallesen, G., Hamilton-Dutoit, S.J., Rowe, M. and Young, L.S. (1991) Expression of Epstein-Barr virus latent gene products in tumour cells of Hodgkin's disease. Lancet. 337, 320-322.

Plebanski, M., Saunders, M., Burtles, S.S., Crowe, S. and Hooper, D.C. (1992). Primary and secondary human *in vitro* T-cell responses to soluble antigens are mediated by subsets bearing different CD45 isoforms. Immunology. 75, 86-91.

Roncarolo, M. and Levings, M.K. (2000). The role of different subsets of T regulatory cells in controlling autoimmunity. *Curr Opin Immunol.* 12, 676-683.

5 Roncarolo, M.G., Levings, M.K., and Traversari, C. (2001). Differentiation of T Regulatory Cells by Immature Dendritic cells. *J. Exp. Med.* 193, F5-F9.

10 Rowe M, Young LS, Crocker J, Stokes H, Henderson S, Rickinson AB. Epstein-Barr virus (EBV)-associated lymphoproliferative disease in the SCID mouse model: implications for the pathogenesis of EBV-positive lymphomas in man. *J Exp Med* 1991;173:147-58.

15 Scheffold A, Assenmacher M, Reiners-Schramm L, Lauster R, Radbruch A. (2000) High-sensitivity immunofluorescence for detection of the pro- and anti-inflammatory cytokines gamma interferon and interleukin-10 on the surface of cytokine-secreting cells. *Nat Med* 6(1):107-10.

20 Schlaak JF, Schmitt E, Huls C, Meyer zum Buschenfelde KH, Fleischer B. (1994) A sensitive and specific bioassay for the detection of human interleukin-10. *J Immunol Methods* 168(1):49-54.

Schleef M, Schmidt T, Flaschel E. (2000) Plasmid DNA for pharmaceutical applications. *Dev Biol (Basel)* 104:25-31

25 Seddon, B., and Mason, D. (2000). The third function of the thymus. *Immunol. Today* 21, 95-99.

Shevach, E.M. (2000). Regulatory T cells in autoimmunity. *Ann. Rev. Immunol.* 18, 423-449.

30 Shevach, E.M., Thornton A., Suri-Payer E., (1998). T lymphocyte-mediated control of autoimmunity. *Novartis Found Symp.* 215, 200-211.

---

Singh, H. and Raghava, G.P.S. (2001) ProPred: Prediction of HLA-DR binding sites. *Bioinformatics*, 17(12), 1236-37.

Smith HA. (2000) Regulation and review of DNA vaccine products. *Dev Biol (Basel)* 104:57-62.

Spencer, J.V., Lockridge, K.M., Barry, P.A., Lin, G., Tsang, M., Penfold, M.E.T., and Schall, T.J. (2002). Potent immunosuppressive activities of Cytomegalovirus encoded Interleukin-10. *J. Virol.* 76, 1285-1292.

5 Stephens, L.A., and Mason, D. (2000). CD25 is a marker for CD4+ thymocytes that prevent autoimmune diabetes in rats, but peripheral T cells with this function are found in both CD25+ and CD25- subpopulations. *J. Immunol.* 165, 3105-3110.

Stott, L.M., Barker, R.N., and Urbaniak, S.J. (2000).  
 10 Identification of alloreactive T-cell epitopes on the Rhesus D protein. *Blood.* 96, 4011-4019.

Sturniolo, T., Bono, E., Ding, J., Raddrizzani, L., Tuereci, O., Sahin, U., Braxenthaler, M., Gallazzi, F., Protti, M.P., Sinigaglia, F., Hammer, J., Generation of tissue-specific and  
 15 promiscuous HLA ligand database using DNA microarrays and virtual HLA class II matrices. *Nat. Biotechnol.* 17, 555-561(1999).

Taylor, P.A., Noelle, R.J., and Blazar, B.R. (2001). CD4+CD25+ immune regulatory cells are required for induction of tolerance to alloantigen via costimulatory blockade. *J. Exp. Med.* 193, 1311-  
 20 1318.

Thomsen, A.R., (2001). The outcome of viral infection is determined by an interplay between CD4+ and CD8+ T cells. *Immunology* 104, (suppl. 1) 40.

Weiner, H.L. (1997). Oral tolerance: Immune mechanisms and  
 25 treatment of autoimmune diseases. *Immunol. Today* 18, 335-343.

Xu, X.-N., Screaton, G.R., and McMichael, A.J. (2001). Virus Infections: Escape, Resistance, and Counterattack. *Immunity* 15, 867-870.

Young, L.S., Eliopoulos, A.G., Gallagher, N.J., and Dawson, C.W.  
 30 (1998). CD40 and epithelial cells: across the great divide. *Immunol. Today* 19, 501-506.

CLAIMS:

1. A method for assessing the tolerogenicity of a test peptide sequence from an infectious agent, comprising the steps of:

(i) contacting a cell population with said test peptide sequence,  
(ii) determining whether IL-10 expression in said cell population is increased, and optionally

(iii) correlating the result of step (ii) with the tolerogenicity of the sequence,

wherein said cell population comprises mononuclear leukocytes from a donor previously infected by said infectious agent.

2. A method according to claim 1, wherein said cell population comprises at least one type of antigen presenting cell.

3. A method according to claim 1 or claim 2, wherein said mononuclear leukocytes comprise at least T lymphocytes, B lymphocytes, natural killer (NK) cells, monocytes, macrophages or dendritic cells.

4. A method according to claim 3, wherein said mononuclear leukocytes comprise at least CD4<sup>+</sup> T lymphocytes.

5. A method according to claim 4, wherein said mononuclear leukocytes further comprise at least one type of antigen presenting cell.

6. A method according to any one of claims 1 to 5, further comprising the steps of:

(i) (a) contacting a similar cell population from a donor not previously infected by said infectious agent with said test peptide sequence; and

(ii) (a) determining whether IL-10 expression in said cell population is increased,

and optionally

(ii) (b) comparing the results from step (ii) with the results from step (ii) (a).

7. A method according to any one of claims 1 to 6, wherein the infectious agent is a virus.

8. A method according to claim 7, wherein the virus is a herpesvirus encoding a viral IL-10 homologue

9. A method according to claim 8, wherein the virus is EBV.

10. A method according to claim 9, wherein the test peptide sequence is derived from EBV LMP1 protein or LMP2 protein.

11. A method according to claim 10, wherein the test or tolerogenic peptide sequence comprises one or more of the sequences P1 to P75 or P1' to P96'.

12. A method for assessing the tolerogenicity of a test peptide sequence from an infectious agent towards a target antigen, comprising the steps of:

(i) contacting a cell population with (a) said test peptide sequence and (b) a target antigen, to make a test composition, and

(ii) re-contacting the cell population from said test composition with said target antigen.

wherein said cell population comprises mononuclear leukocytes from a donor previously infected by said infectious agent.

13. A method according to claim 12, further comprising the steps of:

5 (iii) assessing the response of said cell population to said target antigen, and optionally

(iv) correlating the result of step (iii) with the tolerogenicity of the test peptide sequence.

10 14. A method according to claim 13, wherein step (iii) comprises assessment of cell proliferation or expression of IL-4, IL-2, IL-12 or gamma-IFN.

15 15. A method according to claims 12 to 14, further comprising the step of adding fresh antigen presenting cells prior to step (ii).

20 16. A method according to any one of claims 12 to 15, further comprising the step of contacting the cell population with a confirmatory antigen unrelated to the test sequence or the target antigen.

17. A method according to any one of claims 12 to 16, wherein the infectious agent is a virus.

25 18. A method according to claim 17, wherein the virus is a herpesvirus encoding a viral IL-10 homologue

19. A method according to claim 18, wherein the virus is EBV.

30 20. A method according to claim 19, wherein the test peptide sequence is derived from EBV LMP1 protein or LMP2 protein.

35 21. A method according to claim 20, wherein the test or tolerogenic peptide sequence comprises one or more of the sequences P1 to P75 or P1' to P96'.

22. A method for assessing the tolerogenicity of a test peptide sequence, comprising the steps of:

5 (i) contacting a first cell population with said test peptide sequence,

(ii) contacting a second cell population with a control peptide sequence

0 (iii) determining whether IL-10 expression in each said cell population is increased, and optionally

(iv) correlating the result of step (iii) with the tolerogenicity of the test peptide sequence,

5 wherein each said cell population comprises mononuclear leukocytes from a donor previously infected by an infectious agent, and said control peptide sequence is derived from said infectious agent.

0 23. A method according to claim 22 wherein said control peptide sequence has previously been identified to induce IL-10 expression in a cell population comprising mononuclear leukocytes from a donor previously infected by said infectious agent.

5 24. A method according to claim 21 or claim 22, wherein said first and second cell populations are derived from the same donor.

25. A method according to claim 24, wherein said first and second cell populations comprise a T cell clone capable of proliferating  
0 in response to the control peptide.

26. A method according to any one of claims 22 to 25, wherein said infectious agent is EBV.

5 27. A method according to claim 26, wherein said control peptide is derived from LMF1 or LMF2.

28. A method according to claim 27, wherein said control peptide comprises one or more of P1 to P75 and P1' to P96'.

29. A method of tolerising a cell population to a target antigen,  
comprising contacting said cell population with

(a) a tolerogenic peptide sequence from an infectious agent, or a nucleic acid encoding said test peptide sequence, such that said test peptide sequence is expressed in said cell population; and

(b) the target antigen, or a nucleic acid encoding said test peptide sequence, such that said test peptide sequence is expressed in said cell population;

wherein said cell population comprises mononuclear leukocytes from a subject seropositive for said infectious agent.

30. A method according to claim 29, comprising the steps of contacting a population of antigen presenting cells with said tolerogenic peptide sequence and said target antigen, and subsequently contacting said cell population with said population of antigen presenting cells.

31. A method according to claim 29 or claim 30, wherein said mononuclear leukocytes are contacted with said tolerogenic peptide sequence and said target antigen *in vitro*.

32. A method according to claim 31, wherein said cell population or a subset thereof is re-administered to said subject after contacting with said tolerogenic peptide sequence and said target antigen.

33. A method according to claim 30, wherein said population of antigen presenting cells is contacted with said tolerogenic peptide sequence and said target antigen *in vitro* and said cell population is contacted with said population of antigen presenting cells *in vivo*.



34. A method according to claim 29, wherein said tolerogenic peptide sequence and said target antigen are administered directly to said subject.

35. A method according to any one of claims 29 to 34, wherein the infectious agent is a virus.

36. A method according to claim 35, wherein the virus is a herpesvirus encoding a viral IL-10 homologue.

37. A method according to claim 36, wherein the virus is EBV.

38. A method according to claim 37, wherein the tolerogenic peptide sequence is derived from EBV LMP1 protein or LMP2 protein.

39. A method according to claim 38, wherein the tolerogenic peptide sequence comprises one or more of the sequences P1 to P75 or P1' to P96'.

40. A pharmaceutical composition, for tolerising a subject against a target antigen, comprising a tolerogenic peptide sequence from an infectious agent, or a nucleic acid encoding a tolerogenic peptide sequence from an infectious agent, in admixture with a pharmaceutically acceptable carrier.

41. A pharmaceutical composition according to claim 40, wherein the tolerogenic peptide sequence is derived from EBV.

42. A pharmaceutical composition according to claim 41, wherein the tolerogenic peptide sequence is derived from LMP1 or LMP2.

43. A pharmaceutical composition according to claim 42, wherein the tolerogenic peptide sequence comprises one or more of the peptide sequences P1 to P75 or P1' to P96'.

44. A pharmaceutical composition according to any one of claims 40 to 43, further comprising said target antigen.

5 45. A pharmaceutical composition according to any one of claims 41 to 44, for administration to an individual previously infected with EBV.

10 46. Use of EBV LMP1, LMP2, or a fragment or mimetic thereof, or a nucleic acid encoding the same, in the preparation of a medicament for the prophylaxis or treatment of a condition mediated by an immune response directed against a target antigen.

15 47. Use according to claim 46, wherein the medicament is formulated for administration in conjunction with the target antigen or a nucleic acid encoding the target antigen.

20 48. Use according to claim 46 or 47, wherein the medicament comprises the target antigen or a nucleic acid encoding the target antigen.

25 49. Use according to any one of claims 46 to 48, wherein the condition is type I diabetes mellitus, coeliac disease, multiple sclerosis, rheumatoid arthritis, systemic lupus erythaematosus, myaesthesia gravis, autoimmune haemolytic anaemia, thrombocytopenia, an atopic response or allergy, or a response to a therapeutic product.

30 50. Use according to any one of claims 46 to 48 wherein the target antigen is a cell.

51. Use according to claim 50 wherein the cell is for transplantation.

35 52. Use according to claim 51 in combination with claim 48 wherein the cell comprises nucleic acid encoding the tolerogenic peptide sequence.

53. Use according to any one of claims 46 to 52 wherein the medicament is formulated for administration to an individual previously infected with EBV.

5 54. A peptide having the sequence of any one of P1 to P75 and P1' to P96'.

10 55. A peptide according to claim 54, having the sequence of any one of P2, P4, P5, P6, P7, P8, P9, P10, P12, P13, P14, P15, P16, P17, P18, P20, P22, P23, P24, P25, P26, P27, P29, P30, P32, P34, P35, P39, P68, P71, P72.




Figure 1




Figure 2




Figure 3




Figure 4

BEST AVAILABLE COPY




Figure 5



BEST AVAILABLE COPY




Figure 6




Figure 7a

BEST AVAILABLE COPY




Figure 7b




Figure 7c

BEST AVAILABLE COPY




Figure 7d

BEST AVAILABLE COPY




Figure 8a

BEST AVAILABLE COPY




Figure 8b




Figure 8c






Figure 9a

BEST AVAILABLE COPY




Figure 9b




Figure 9c

BEST AVAILABLE COPY




Figure 9d




Figure 10a

BEST AVAILABLE COPY




Figure 10b




Figure 10c